

BLADE™

SELECTIVE HERBICIDE for TURFGRASS

Control bahiagrass, ryegrass, foxtail and many broadleaf weeds in warm-season turf, as well as Kentucky bluegrass and fine fescue.

Blade™ effectively controls bahiagrass in established bermudagrass and St. Augustinegrass areas. Low use rates also selectively control many broadleaf weeds and other grasses including foxtail and ryegrass.

Blade also controls a wide range of broadleaf weeds including spurge (prostrate), wild onion, wild garlic, clover, dandelion, dollarweed, plantains, chickweed, clover, lespeדה, Florida pusley and many more listed weeds.

Use Blade to remove overseeded ryegrass and hasten spring release of bermudagrass – without injury to the bermudagrass. Blade does not delay the transition to bermudagrass in the spring. Transition occurs slowly over 3 to 4 weeks

Blade is also used in lawns for control of broadleaf weeds in St. Augustinegrass turf. Blade is effective at very low rates, ranging from 0.125 to 1.0 ounce of product per acre, yet has a good toxicity profile for animals and humans.


Use Blade only on bermudagrass, St. Augustinegrass, zoysiagrass (Meyer and Emerald), centipedegrass, Kentucky bluegrass and fine fescue (excluding tall fescue and turf-type tall fescue). Turf species have a high degree of tolerance at labeled use rates.

Blade is a trademark of PBI-GORDON CORPORATION.


QuickSilver is a registered trademark of FMC Corporation.




KNOTWEED



WILD GARLIC



WILD ONIONS



DOLLARWEED



BAHIAGRASS



BERMUDA RELEASE

- CAUTION signal word
- Very low use rates at very affordable cost
- Hastens spring release of overseeded bermudagrass
- Transition occurs slowly over 3 to 4 weeks
- Overseed in 8 weeks
- Rain-fast in 6-8 hours

- Controls ryegrass, bahiagrass and foxtail
- Controls broadleaf weeds such as spurge, wild onion, wild garlic, dandelion, dollarweed and many more
- For best results, use a non-ionic surfactant
- May be tank-mixed with other herbicides, such as QuickSilver®



Contains
metsulfuron
methyl

Extruded Granules

RECOMMENDATIONS

Post-emergence applications at low rates to control bahiagrass, many broadleaf weeds and certain grasses in warm-season turf, as well as Kentucky bluegrass and fine fescue turf. Sites include golf course (fairways, aprons, tees and roughs), lawns, parks, cemeteries and other similar non-crop areas. This product may also be used on sod farms.

ACTIVE INGREDIENTS:

Metsulfuron Methyl	
Methyl 2-[[[(4-methoxy-6-methyl-1,3,5-triazin-2-yl)amino]carbonyl]amino]sulfonyl]benzoate	60%
Inert Ingredients	40%

EPA Reg. No. 66222-164-2217

PACKAGING:

Plastic 4 X 2 oz. with measuring device
per case: shipping weight 2 lbs.

READ LABEL BEFORE USE

The literature contained herein is not intended to be used as a substitute for the information contained on the label of the product container. Specimen labels and other literature are subject to revision. Before using this product, read and follow all label instructions on the container/package.

pbigordon.com/blade

weedalert.com



An Employee-Owned Company

1217 W. 12TH STREET • P.O. BOX 014090
KANSAS CITY, MISSOURI 64101-0090
816.421.4070 • FAX 816.474.0462
800.821.7925

© 2010, PBI-Gordon Corporation