

CLICK HERE TO JUMP TO USE DIRECTIONS



# STUMP KILLER READY-TO-USE

KILLS STUMPS DOWN TO THE ROOTS

- PREVENTS RE-GROWTH
- KILLS EXISTING SPROUTS
- CONVENIENT SQUIRT BOTTLE
- APPLY DIRECTLY ON STUMP

**ACTIVE INGREDIENT:**

Glyphosate (N-(phosphonomethyl) glycine) in the form of its isopropylamine salt . . . . . 32.3%

**INERT INGREDIENTS:** . . . . . 67.7%  
TOTAL 100.0%

**THIS PRODUCT CONTAINS:**

3 lbs. per U.S. gallon of the active ingredient glyphosate in the form of its isopropylamine salt.

**KEEP OUT OF REACH OF CHILDREN  
CAUTION**



**READ THE ENTIRE LABEL FIRST.  
OBSERVE ALL PRECAUTIONS AND  
FOLLOW DIRECTIONS CAREFULLY.**

**PRECAUTIONARY STATEMENTS**

**Hazards to Humans and Domestic Animals**

**CAUTION:** Causes moderate eye irritation. Harmful if absorbed through the skin. Harmful if swallowed. Avoid contact with skin, eyes or clothing. Wash thoroughly with soap and water after handling.

**Personal Protective Equipment (PPE)**

Some materials that are chemical resistant to this product are listed below. If you want more options, follow instructions for Category A on an EPA chemical resistant category selection. Applicators and other handlers must wear: long-sleeved shirt and long pants, chemical resistant gloves Category A, such as butyl rubber, natural rubber, neoprene rubber, or nitrile rubber ≥ 14 mils, and shoes plus socks.

**Engineering Control Statements**

When handlers use enclosed cabs or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240(d)(4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

**User Safety Recommendations**

- Users should wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- Users should remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Users should remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

**First Aid**

- If swallowed:**
- Call a poison control center or doctor immediately for treatment advice.
  - Have person sip a glass of water if able to swallow.
  - Do not induce vomiting unless told to do so by the poison control center or doctor.
  - Do not give anything by mouth to an unconscious person.

(cont. on next column)

First Aid (cont.)	
<b>If on skin or clothing:</b>	<ul style="list-style-type: none"> <li>• Take off contaminated clothing.</li> <li>• Rinse skin immediately with plenty of water for 15-20 minutes.</li> <li>• Call a poison control center or doctor for treatment advice.</li> </ul>
<b>If inhaled:</b>	<ul style="list-style-type: none"> <li>• Move person to fresh air.</li> <li>• If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible.</li> <li>• Call a poison control center or doctor for treatment advice.</li> </ul>
<b>If in eyes:</b>	<ul style="list-style-type: none"> <li>• Hold eye open and rinse slowly and gently with water for 15-20 minutes.</li> <li>• Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.</li> <li>• Call a poison control center or doctor for treatment advice.</li> </ul>
Have the product container or label with you when calling a poison control center or doctor or going for treatment.	

**Domestic Animals**

This product is considered to be relatively nontoxic to dogs and other domestic animals; however, ingestion of this product or large amounts of freshly sprayed vegetation may result in temporary gastrointestinal irritation (vomiting, diarrhea, colic, etc.) If such symptoms are observed, provide the animal with plenty of fluids to prevent dehydration. Call a veterinarian if symptoms persist for more than 24 hours.

**Environmental Hazards**

Do not contaminate water when disposing of equipment washwaters and rinsate. Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwaters and rinsate.

**Physical or Chemical Hazards**

Combustible. Do not use or store near heat or open flame.

Spray solutions of this product should be mixed, stored and applied using only stainless steel, aluminum, fiberglass, plastic or plastic-lined steel containers.

DO NOT MIX, STORE OR APPLY THIS PRODUCT OR SPRAY SOLUTIONS OF THIS PRODUCT IN GALVANIZED STEEL OR UNLINED STEEL (EXCEPT STAINLESS STEEL) CONTAINERS OR SPRAY TANKS. This product or spray solutions of this product react with such containers and tanks to produce hydrogen gas which may form a highly combustible gas mixture. This gas mixture could flash or explode causing serious personal injury, if ignited by open flame, spark, welder's torch, lighted cigarette or other ignition source.

**DIRECTIONS FOR USE**

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide registration.

**Agricultural Use Requirements**

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170.

This standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard. Do not enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 4 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil or water is: coveralls, waterproof gloves, and shoes plus socks.

**Non-Agricultural Use Requirements**

The requirements in this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard (40 CFR Part 170) for agricultural pesticides. The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries, or greenhouses.

Keep people and pets off treated areas until spray solution has dried.

**Cut-Surface Stump Application**

Woody vegetation may be controlled by treating freshly cut stumps of trees and resprouts with this product. Cut stump treatments are effective on stumps with diameters larger than 3 to 4 inches in diameter. Apply this product using suitable equipment to ensure coverage of the entire cambium. Apply Stump Killer Ready-To-Use undiluted directly to freshly cut surface immediately after cutting. Delay in applying this product may result in reduced performance. For best results, trees should be cut during periods of active growth and full leaf expansion.

- Cut trees and brush close to the soil surface. Apply this product to the freshly cut surface of the stump *immediately* after cutting. Delays in application may reduce the effectiveness.
- Apply for full and thorough coverage of the cut surface. Treat the entire circumference of the stump by drenching the cambium (the outermost wood between the pith and the bark).
- Treatment will control or suppress many types of woody brush and trees, such as alder, bluegum eucalyptus, madrone, oak, giant reed, saltcedar, sweetgum and tan oak.
- The most favorable period for cut stump treatment occurs after full leaf stage in the spring and continues into early summer. Brush and trees treated in the late fall may not be fully controlled.
- Follow-up applications of this product or with a different herbicide may be needed under less favorable growing conditions and for hard-to-control species. Another choice for effective control with a cut-surface stump treatment is Gordon's Cleanout® Brush and Stump Spray.

When used according to directions for cut stump application, this product will control, partially control or suppress most woody brush and tree species, some of which are listed below:

Alder <i>Alnus spp.</i>	Oak <i>Quercus spp.</i>
Coyote brush <i>Baccharis consanfuinea*</i>	Poplar <i>Populus spp.*</i>
Dogwood <i>Cornus spp.*</i>	Reed, giant <i>Arundo donax</i>
Eucalyptus <i>Eucalyptus spp.</i>	Saltcedar <i>Tamarix spp.</i>
Hickory <i>Carya spp.*</i>	Sweetgum <i>Liquidambar styraciflua*</i>
Madrone <i>Arbutus menziesii</i>	Sycamore <i>Platanus occidentalis*</i>
Maple <i>Acer spp.*</i>	Tanoak <i>Lithocarpus densiflorus</i>
	Willow <i>Salix spp.</i>

\*This product is not approved for this use on these species in the State of California.

## For Brush Control

### Foliar Sprays for Brush

- Cut back brush that is taller than 5 feet and then spray regrowth.
- For maximum activity, brush should be growing vigorously at the time of applications. The most favorable period for treatment occurs after full leaf stage in the spring and continues into early summer. Brush treated in the late fall may not be fully controlled.
- Spray uniformly the foliage of the vegetation to be controlled. Apply for full coverage and thorough brush/weed contact.
- Follow-up applications of this product or with a different herbicide may be needed under less favorable growing conditions and for hard-to-control species including blackberry, poison oak, and kudzu. Another choice for effective brush control is Gordon's Cleanout® Brush and Stump Spray. Cleanout® is recommended to control undesirable woody plants including buckbrush, poison ivy, multiflora rose, and sumac. Foliar applications of Cleanout® are appropriate during the period of active growth of the brush. Cut stump, basal bark, and frill treatments of Gordon's Cleanout® Brush and Stump Spray may be applied any time of year.

### Recommended Use Rates and Species Controlled

Stump Killer Ready-To-Use may be used as a broadcast treatment or spot treatment with any of the application equipment listed on this label in noncropland sites.

### Woody Brush and Trees

Handy-held high volume sprayer: Amount of product to add per 1 gallon of spray solution for spray-to-wet coverage is 11.5 fl.oz.

### Species:

Alder <i>Alnus spp.</i>	Hasardia <i>Haplopappus squamosus*</i>
Ash <i>Fraxinus spp.*</i>	Hawthorn <i>Crataegus spp.</i>
Aspen, quaking <i>Populus tremuloides</i>	Hazel <i>Corylus spp.</i>
Bearmat (Bearclover) <i>Chamaebatia foliolosa</i>	Hickory <i>Carya spp.</i>
Beech	Holly, Florida/Brazilian peppertree
Birch <i>Betula spp.</i>	<i>Schinus terebinthifolius</i>
Blackberry <i>Rubus spp.</i>	Honeysuckle <i>Lonicera spp.</i>
Blackgum	Hornbeam, American <i>Carpinus caroliniana</i>
Bracken	Kudzu <i>Pueraria lobata</i>
Broom: French <i>Cytisus monspessulanus</i>	Locust, black <i>Robinia pseudoacacia*</i>
Scotch <i>Cytisus scopadus</i>	Madrone resprouts
Buckwheat, California	Manzanita <i>Arctostaphylos spp.</i>
<i>Edogonum fasciculatum*</i>	Maple: Red <i>Acer rubrum*</i>
Cascara <i>Rhamnus purshiana*</i>	Sugar <i>Acer saccharum</i>
Catsclaw <i>Acacia greggii*</i>	Vine <i>Acer circinatum*</i>
Ceanothus <i>Ceanothus spp.</i>	Monkey Flower <i>Mimulus guttatus*</i>
Chamise <i>Adenostoma fasciculatum</i>	Oak: Black <i>Quercus yelutina*</i>
Cherry: Bitter <i>Prunus emarginata</i>	Northern Pin <i>Quercus palustris</i>
Black <i>Prunus serotina</i>	Post <i>Quercus stellata</i>
Pin <i>Prunus pensylvanica</i>	Red <i>Quercus rubra</i>
Coyotebrush <i>Bacchads consanguinea</i>	Southern Red <i>Quercus falcata</i>
Creeper, Virginia <i>Parthenocissus quiflquefoja*</i>	White <i>Quercus alba*</i>
Dewberry <i>Rubus trivialis</i>	Persimmon <i>Diospyros spp.*</i>
Dogwood <i>Cornus spp.</i>	Pine
Elderberry <i>Sambucus spp.</i>	Poison ivy <i>Rhus radicans</i>
Elm <i>Ulmus spp.*</i>	Poison oak <i>Rhus toxicodendron</i>
Eucalyptus, bluegum <i>Eucalyptus spp.</i>	
Gorse	

(cont. on next column)

### Species (cont.):

Poplar, yellow (Tulip tree) <i>Liriodendron tulipifera*</i>	Sourwood <i>Oxydendrum arboreum*</i>
Prunus <i>Prunus spp.†</i>	Sumac: Poison <i>Rhus vernix*</i>
Rasperry <i>Rubus spp.</i>	Smooth <i>Rhus glabra*</i>
Redbud, eastern <i>Cercis canadensis</i>	Winged <i>Rhus copallina*</i>
Rose, multiflora <i>Rosa multiflora</i>	Sweetgum <i>Liquidambar styraciflua</i>
Russian olive <i>Elaeagnus angustifolia</i>	Swordfern <i>Polystichum munitum</i>
Sage, black, <i>white Salvia spp.</i>	Tallowtree, Chinese <i>Sapium sebiferum</i>
Sagebrush, California <i>Artemisia californica</i>	Thimbleberry <i>Rubus parviflorus</i>
Salmonberry <i>Rubus spectabilis</i>	Tobacco, tree <i>Nicotiana glauca*</i>
Saltcedar <i>Tamarix spp.</i>	Toyon
Saltbush, sea myrtle <i>Baccharis halimifolia</i>	Trumpetreeper <i>Campsis radicans</i>
Sassafras <i>Sassafras spp.</i>	Waxmyrtle, southern <i>Myrica cerifera*</i>
	Willow <i>Salix spp.</i>
	Yerbasanta

† Not for use in California

\*Partial control

**NOTE:** If brush has been mowed or tilled or trees have been cut, do not treat until regrowth has occurred. Apply this product when plants are actively growing and after full leaf expansion. Best results are obtained when application is made in late summer or fall after fruit formation. In arid areas, best results are obtained when application is made in the spring to early summer when brush species are at high moisture content and are flowering. Ensure thorough coverage when using hand-held equipment. Symptoms may not appear prior to frost or senescence with fall treatments. Allow 7 or more days after application before tillage, mowing or removal. Repeat treatments may be necessary to control plants regenerating from underground parts or seed. Some autumn colors on undesirable deciduous species are acceptable provided no major leaf drop has occurred. Reduced performance may result if fall treatments are made following a frost.

## General Information

- This product mixes readily with water and is intended to be applied as a foliar spray for the control or destruction of many herbaceous or woody plants. It may be applied through most standard industrial or field-type sprayers after dilution and thorough mixing with water in accordance with label instructions.
- This product moves through the plant from the point of foliage contact to and into the root system. Visible effects on most annual weeds occur within 2 to 4 days, but on most perennial weeds may not occur for 7 days or more. Extremely cool or cloudy weather following treatment may slow activity of this product and delay visual effects of control.
- Visible effects are gradual wilting and yellowing of the plant which advances to complete browning of above-ground growth and deterioration of underground plant parts. Unless otherwise specified on this label, delay application until vegetation has emerged and reached the growth stages described for control. Unemerged plants arising from unattached underground rhizomes or root stocks of perennials will not be affected by the herbicide and will continue to grow. For this reason, best control of most perennial weeds is obtained when treatment is made at late growth stages approaching maturity.
- Always use the higher rate of this product when the weed growth is heavy or dense.
- Reduced control may result when applications are made to annual or perennial weeds that have been mowed, grazed, or cut, and have not been allowed to re-grow to the recommended stage for treatment. Repeat treatments will be necessary to control the growth of later germinating seeds.
- Rainfall or irrigation occurring within 6 hours after application may reduce effectiveness. Heavy rainfall or irrigation within 2 hours after application may wash the chemical off the foliage and a repeat treatment may be required.
- This product does not provide residual weed control. For subsequent residual weed control, follow a label-approved herbicide program. Read and carefully observe the cautionary statements and all other information appearing on the labels of all herbicides used.
- Buyer and all users are responsible for all loss or damage in connection with the use or handling of mixtures of this product with herbicides or other materials that are not expressly recommended in this labeling. Mixing this product with herbicides or other materials not recommended on this label may result in reduced performance.
- For best results, spray coverage should be uniform and complete.

### Use Precautions

- AVOID DRIFT. EXTREME CARE MUST BE USED WHEN APPLYING THIS PRODUCT TO PREVENT INJURY TO DESIRABLE PLANTS AND CROPS.
- Avoid treating weeds under poor growing conditions such as drought stress, disease or insect damage, as reduced weed control may result. Reduced results may also occur when treating weeds heavily covered with dust.
- Do not allow the herbicide solution to mist, drip, drift or splash onto desirable vegetation since minute quantities of this product can cause severe damage or destruction to the crop, plants, or other areas on which treatment was not intended.
- The likelihood of injury occurring from the use of this product is greatest when winds are gusty or in excess of 5 miles per hour or when other conditions, including lesser wind velocities, will allow spray drift to occur. When spraying, avoid combinations of pressure and nozzle types that will result in splatter or fine particles (mist) which are likely to drift.
- AVOID APPLYING AT EXCESSIVE SPEED OR PRESSURE.
- NOTE: Use of this product in any manner not consistent with this label may result in injury to persons, animals or crops, or other unintended consequences. Keep container closed to prevent spills and contamination.

## Spray Preparation and Tank Mixes

Stump Killer Ready-To-Use is a soluble concentrate that dissolves readily in clean water (free of mud or clay). Reduced results may occur if water containing soil is used, such as water from ponds or unlined ditches.

Additional surfactants are not required for this formulation.

### Stump Killer Ready-To-Use Alone With Water As Diluent

1. Determine spray volume needed for the treated area. Add one-half the required amount of clean water to the spray tank.
2. Measure the recommended amount of product and add Stump Killer Ready-To-Use slowly with agitation, and complete filling the tank with water.
3. Mix thoroughly and continue agitation while spraying.

### Colorants or Dyes

Colorants or marking dyes may be added to this product. Colorants or dyes used in spray solutions of this product may reduce performance, especially at lower rates or dilutions. Use colorants or dyes according to the manufacturer's recommendations.

### Drift Control Additives

Drift control additives may be added to this product. When a drift control additive is used, read and carefully observe the cautionary statements and all other information appearing on the additive label.

### Sprayer Cleanup

CLEAN THE ENTIRE SPRAYER AFTER APPLICATION OF THIS PRODUCT. Rinse sprayer and all parts with water three (3) times. Spray rinse water on bare soil or gravel. After cleaning, sprayer may be used for other products. Failure to clean the sprayer thoroughly may result in injury to desirable crops which are subsequently sprayed.

## Application Equipment and Techniques

Do not apply this product through any type of irrigation system.

This product may be applied with the following application equipment:

- **Hand-held low-volume and high-volume spray equipment** - knapsack and backpack sprayers, pump-up pressure sprayers, spray guns, hand wands, lances and other hand-held and motorized spray equipment used to direct the spray onto weed foliage. Applications should be made on a spray-to-wet basis; spray coverage should be uniform and complete. Use only coarse sprays. If a straight-stream nozzle is used, start the application at the top of the target vegetation and spray from top to bottom in a "zigzag" motion. Ensure that the spray contacts 50% or more of the leaf surface. Small open-branched trees need only be treated from one side. If foliage is thick or there are multiple root sprouts, application must be made from several sides to ensure adequate coverage. When using application methods that result in less than complete coverage, use the higher recommended rate.

### EQUIVALENT MEASUREMENTS:

- 1 Tablespoon (Tbs.) = 3 teaspoon (tsp.)
- 1 fluid ounce (fl.oz.) = 2 Tablespoons (Tbs.)
- 2 fluid ounces (fl.oz.) = 4 Tablespoons (Tbs.)

## STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

**PESTICIDE STORAGE:** Store in original container in a locked storage area inaccessible to children or pets. Keep from freezing.

**PESTICIDE DISPOSAL:** Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

**CONTAINER HANDLING:** Nonrefillable container. Do not reuse or refill this container. Offer for recycling, if available, or puncture and dispose of in a sanitary landfill, or by incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

Triple rinse or pressure rinse container (or equivalent) promptly after emptying.

Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times.

OR

Pressure rinse as follows: Empty the remaining contents into application equipment or a mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank or collect rinsate for later use or disposal. Insert pressure rinsing nozzle in the side of the container, and rinse at about 40 PSI for at least 30 seconds. Drain for 10 seconds after the flow begins to drip.

## LIMITED WARRANTY AND DISCLAIMER

The manufacturer warrants only that the chemical composition of this product conforms to the ingredient statement given on the label, and that the product is reasonably suited for the labeled use when applied according to the Directions for Use.

THE MANUFACTURER NEITHER MAKES NOR INTENDS ANY OTHER EXPRESS OR IMPLIED WARRANTIES, INCLUDING ANY WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE, WHICH ARE EXPRESSLY DISCLAIMED. This limited warranty does not extend to the use of the product inconsistent with label instructions, warnings or cautions, or to

use of the product under abnormal conditions such as drought, excessive rainfall, tornadoes, hurricanes, etc. These factors are beyond the control of the manufacturer or the seller. Any damages arising from a breach of the manufacturer's warranty shall be limited to direct damages, and shall not include indirect or consequential damages such as loss of profits or values, except as otherwise provided by law.

The terms of this Limited Warranty and Disclaimer cannot be varied by any written or verbal statements or agreements. No employee or agent of the seller is authorized to vary or exceed the terms of this Limited Warranty and Disclaimer in any manner.

Cleanout® is a registered trademark of PBI/Gordon Corporation.

955/12-2010 AP061009  
EPA REG. NO. 2217-852



An Employee-Owned Company

MANUFACTURED BY  
PBI/GORDON CORPORATION  
1217 WEST 12TH STREET  
KANSAS CITY, MISSOURI 64101  
[www.pbigordon.com](http://www.pbigordon.com)