

[CLICK HERE TO JUMP TO USE DIRECTIONS](#)



SYNERGIZED LICE-NO-MORE™ INSECTICIDE

For Cattle, Sheep* and Their Premises
Synergized Formula For Effective Control
Can Be Used on Lactating Dairy Cattle

Ready-To-Use as a Pour-On, Spray or Mist –
or mix for Backrubber use.

Also Kills Flies and Mosquitoes

ACTIVE INGREDIENTS:

Permethrin [^{**} (3-Phenoxyphenyl) methyl (+ or -) cis-trans-3-(2,2-dichloroethenyl) 2,2-dimethylcyclopropanecarboxylate]	1.00%
^{***} Piperonyl Butoxide Technical	1.00%
^{****} OTHER INGREDIENTS	98.00%
TOTAL 100.00%	

^{**} Cis-trans isomers ratio: Min. 35% (+ or -) cis and Max. 65% (+ or -) trans.
^{***} Equivalent to Min. 0.8% (butylcarbityl)(6-propylpiperonyl) ether and 0.2% related compounds.
^{****} Contains petroleum distillate.

**KEEP OUT OF REACH OF CHILDREN
CAUTION**



**READ THE ENTIRE LABEL FIRST.
OBSERVE ALL PRECAUTIONS AND
FOLLOW DIRECTIONS CAREFULLY.**

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION: Harmful if absorbed through skin or inhaled. Avoid contact with skin, eyes and clothing. Avoid breathing vapor or spray mist. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, or using tobacco. Remove and wash contaminated clothing before reuse.

First Aid	
IF SWALLOWED:	<ul style="list-style-type: none"> Immediately call a poison control center or doctor. Do not induce vomiting unless told to do so by a poison control center or doctor. Do not give any liquid to the person. Do not give anything by mouth to an unconscious person.
IF ON SKIN OR CLOTHING:	<ul style="list-style-type: none"> Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.
IF INHALED:	<ul style="list-style-type: none"> Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible. <p style="text-align: right;"><i>(cont. on next column)</i></p>

First Aid (cont.)	
IF INHALED (cont.):	• Call a poison control center or doctor for further treatment.
NOTE TO PHYSICIAN: Contains petroleum distillate – vomiting may cause aspiration pneumonia hazard. Have the product container or label with you when calling a poison control center or doctor, or going for treatment. For information regarding medical emergencies or pesticide incidents, call 1-877-800-5556 .	

ENVIRONMENTAL HAZARDS

This product is extremely toxic to fish and other aquatic invertebrates. Do not add directly to water. Do not contaminate water when disposing of equipment washwaters. Apply the product only as specified on the label.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

READY TO USE - No dilution necessary.

Apply To	Target Insects	Application Instructions
Lactating and Non-lactating Dairy Cattle, Beef Cattle and Calves	Lice, Horn Flies, Face Flies, Aids in control of Horse Flies, Stable Flies, House Flies, Mosquitoes and Black Flies.	DOSEAGE: Apply ½ fl. oz. (15cc) per 100 lbs. body weight of animal up to a maximum of 5 fl. oz. for any one animal. POUR ON: Pour correct dose along back and down face. READY-TO-USE SPRAY: Use undiluted in a mist sprayer to apply correct dose. Apply directly to neck, face, back, legs and ears. BACKRUBBER USE: Mix one pint per gallon #2 diesel or mineral oil. Keep rubbing device charged. Results improved by daily forced use.
Sheep*	Sheep Keds*, Lice*.	Pour along back. Apply ¼ fl. oz. (7.5cc) per 50 lbs. body weight of animal, up to a maximum of 3 fl. oz. for any one animal. For optimum control, all animals in the flock should be treated after shearing.
Premises – in and around Horse, Beef, Dairy, Swine, Sheep and Poultry premises. Animal Hospital Pens and Kennels and "outside" Meat Processing Premises.	House Flies, Stable Flies, Face Flies, Gnats, Mosquitoes, Black Flies, Fleas, Little House Flies (<i>Fannia spp</i>). Aid in control of cockroaches, ants, spiders and crickets.	For use as a ready-to-use spot spray or premise spray, use undiluted in a mist sprayer. Apply directly to surface to leave a residual insecticidal coating, paying particular attention to areas where insects crawl or alight. One gallon will treat approximately 7,300 square feet.

APPLICATION INSTRUCTIONS:

Apply ½ fl. oz. per 100 lbs. body weight of animal, up to a maximum of 5 fl. oz. for any one animal to kill Lice, Horn Flies, and Face Flies. Aids in the insecticidal control of Horse Flies, Stable Flies, Mosquitoes and Black Flies on lactating and non-lactating dairy cattle, beef cattle, calves and sheep.

For sheep, apply ¼ fl. oz per 50 lbs. body weight of animal, up to a maximum of 3 fl. oz for any one animal. Pour-On: Pour along back line and down face. For Keds* on sheep*, pour along back line over neck, shoulders and rump.

Ready-To-Use Spray: Use undiluted in a mist sprayer to apply correct dose. Apply directly to neck, face, back, legs and ears.

For Pour-On and Ready-To-Use Spray applications, repeat treatment as needed but do not exceed 5 fl. oz. for any one animal on cattle or 3 fl. oz. for any one animal on sheep during any 14 day interval. For optimum insecticidal effectiveness against lice, a second treatment at 14 day intervals is recommended.

Backrubber Use: Mix one pint per gallon #2 diesel or mineral oil. Keep rubbing device charged. For Backrubber applications, results are improved by daily forced use.

For cattle and sheep, repeat treatment as needed, but not more often than once every two weeks. For optimum lice control, two treatments at a 14-day interval are recommended.

SPECIAL NOTE: SYNERGIZED LICE-NO-MORE™ INSECTICIDE is not effective in controlling cattle grubs.

SYNERGIZED LICE-NO-MORE INSECTICIDE is an oil-based, ready-to-use product, that may leave an oily appearance on the hair coat of some animals.

SYNERGIZED LICE-NO-MORE INSECTICIDE should be used in an integrated pest management system which may involve repeat treatments and the use of other pest control practices.

*Not registered in the State of California for the control of Sheep Keds and Lice on Sheep.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage and disposal.

PESTICIDE STORAGE: Store in a cool, dry area. Always store pesticides in the original container. Store away from food, pet food and feed.

PESTICIDE DISPOSAL: To avoid wastes, use all material in this container by application according to label directions. If wastes cannot be avoided, offer remaining product to a waste disposal facility or pesticide disposal program (often such programs are run by state or local governments or by industry).

CONTAINER HANDLING: Nonrefillable container. Do not reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Then offer for recycling if available, or puncture and dispose of in a sanitary landfill or by incineration. Do not burn unless allowed by state and local ordinances.

LICE-NO-MORE® is a trademark of PBI-Gordon Corporation.

729/5-2010 AP051409
EPA REG. NO. 1021-1739-2217



An Employee-Owned Company

DISTRIBUTED BY
PBI/GORDON CORPORATION
1217 WEST 12TH STREET
KANSAS CITY, MISSOURI 64101
www.pbigordon.com