

CLICK HERE TO JUMP TO USE DIRECTIONS



# GOAT & SHEEP SPRAY READY-TO-USE

*Insect control for goats, sheep  
and other FARM ANIMALS.*

*A Ready-To-Use Stable Emulsion Insecticide.*

**ACTIVE INGREDIENT:**

\*Permethrin ..... 0.50%

**INERT INGREDIENTS:** ..... 99.50%

TOTAL 100.00%

\*(3-phenoxyphenyl) methyl (+/-) cis/trans 3-(2,2-dichloroethyl)-  
2,2-dimethylcyclopropanecarboxylate. Cis/trans ratio: min. 35% (+/-) cis  
and max. 65% (+/-) trans.

## KEEP OUT OF REACH OF CHILDREN CAUTION

*See below for additional Precautionary Statements and First Aid.*

*Shake well before using.*



**READ THE ENTIRE LABEL FIRST.  
OBSERVE ALL PRECAUTIONS AND  
FOLLOW DIRECTIONS CAREFULLY.**

### PRECAUTIONARY STATEMENTS

**Hazards to Humans & Domestic Animals**

**CAUTION:** Wash thoroughly after handling and before smoking or eating. Avoid contamination of feed and foodstuffs. Remove pets and birds and cover fish aquaria before surface applications. Do not use on humans. Do not allow children or pets to contact treated areas until surfaces are dry.

| FIRST AID  |  |
|--|--|
| <b>If inhaled:</b>   | <ul style="list-style-type: none"> <li>• Move person to fresh air.</li> <li>• If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible.</li> <li>• Call a poison control center or doctor for treatment advice.</li> </ul>                    |
| <b>If on skin or clothing:</b>   | <ul style="list-style-type: none"> <li>• Take off contaminated clothing.</li> <li>• Rinse skin immediately with plenty of water for 15-20 minutes.</li> <li>• Call a poison control center or doctor for treatment advice.</li> </ul>  |
| <b>If in eyes:</b>   | <ul style="list-style-type: none"> <li>• Hold eye open and rinse slowly and gently with water for 15-20 minutes.</li> <li>• Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.</li> <li>• Call a poison control center or doctor for treatment advice.</li> </ul>            |
| <b>If swallowed:</b>   | <ul style="list-style-type: none"> <li>• Immediately call a poison control center or doctor.</li> <li>• Do not induce vomiting unless told to by a poison control center or doctor.</li> <li>• Do not give any liquid to the person.</li> <li>• Do not give anything by mouth to an unconscious person.</li> </ul> |
| <p>Have the product container or label with you when calling a poison control center or doctor or going for treatment. For additional information in case of emergency, call toll free 1-877-800-5556.</p> |  |

**ENVIRONMENTAL HAZARDS:**

This product is toxic to fish. Do not apply directly to water, to areas where surface water is present, or to intertidal areas below the high water mark. Do not contaminate water when disposing of equipment washwater.

**PHYSICAL OR CHEMICAL HAZARDS:**

Do not apply this product in or on electrical equipment due to the possibility of shock hazard.

### DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

### STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

**PESTICIDE STORAGE:** Store upright at room temperature. Avoid exposure to extreme temperatures.

**PESTICIDE DISPOSAL:** Pesticide or rinse water that cannot be used according to label instructions must be disposed of on site or at or by an approved waste disposal facility.

**CONTAINER DISPOSAL:** Do not reuse container. Wrap container in several layers of newspaper and discard in trash.

Do not transfer to another container.

### SHAKE WELL BEFORE USING

### ON GOATS, SHEEP, CATTLE (Beef and Dairy), HOGS and HORSES

**To Control Horn Flies, Face Flies, House Flies, Mosquitoes and Gnats:** Apply a light mist sufficient to wet the surface of the hair. For face flies, spray face and head, but do not spray into eyes.

**To Control Stable Flies, Horse Flies and Deer Flies:** Apply 2 fluid ounces per adult animal, sufficient to wet the hair thoroughly. Repeat treatment daily.

**To Control Blood-Sucking Lice:** Apply to the infested areas of the animal, using a stiff brush to get the spray to the base of the hair. Repeat every 3 weeks.

### POULTRY AND POULTRY HOUSES

**To Control Poultry Lice:** Spray roosts, walls and nests or cages thoroughly. This should be followed by spraying over the birds with a fine mist.

**To Control Bedbugs and Mites in Poultry Houses:** Spray crevices of roost poles, cracks in walls, and cracks in nests where the bedbugs and mites hide.

### LIMITED WARRANTY AND DISCLAIMER

The manufacturer warrants only that the chemical composition of this product conforms to the ingredient statement given on the label, and that the product is reasonably suited for the labeled use when applied according to the Directions for Use.

**THE MANUFACTURER NEITHER MAKES NOR INTENDS ANY OTHER EXPRESS OR IMPLIED WARRANTIES, INCLUDING ANY WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE, WHICH ARE EXPRESSLY DISCLAIMED.** This limited warranty does not extend to the use of the product inconsistent with label instructions, warnings or cautions. Any damages arising from a breach of the manufacturer's warranty shall be limited to direct damages, and shall not include indirect or consequential damages such as loss of profits or values, except as otherwise provided by law.

The terms of this Limited Warranty and Disclaimer cannot be varied by any written or verbal statements or agreements. No employee or agent of the seller is authorized to vary or exceed the terms of this Limited Warranty and Disclaimer in any manner.

763/1-2004 AP031803  
EPA REG. NO. 2217-825

Manufactured For  
 **pbi/GORDON**  
**CORPORATION**  
An Employee-Owned Company  
1217 West 12th Street  
Kansas City, Missouri 64101  
[www.pbigordon.com](http://www.pbigordon.com)