

CLICK HERE TO JUMP TO USE DIRECTIONS

GORDON'S®

LIQUID FRUIT TREE SPRAY

Protects listed fruit and strawberries and roses, evergreens and flowers from both insects and disease.



ACTIVE INGREDIENTS:

Captan (CAS No. 133-06-2)	11.76%
Related derivatives	0.24%
Malathion (CAS No. 121-75-5)	6.00%
Carbaryl (CAS No. 63-25-2)	0.30%
OTHER INGREDIENTS:	81.70%
TOTAL 100.00%	

Contains Petroleum Distillate

KEEP OUT OF REACH OF CHILDREN WARNING

See below for additional Precautionary Statements and First Aid.



**READ THE ENTIRE LABEL FIRST.
OBSERVE ALL PRECAUTIONS AND
FOLLOW DIRECTIONS CAREFULLY.**

PRECAUTIONARY STATEMENTS

Hazards to Humans & Domestic Animals

WARNING: Causes substantial but temporary eye injury or skin irritation. Harmful if swallowed or inhaled. Do not get in eyes, on skin, or on clothing. Avoid breathing spray mist. Wear coveralls worn over short-sleeved shirt and short pants, socks and chemical resistant footwear, goggles or face shield and chemical resistant gloves (such as Barrier Laminate or Viton, Selection Category G). Overhead Exposure: Wear chemical resistant headgear. For Cleaning Equipment: Add a chemical resistant apron. Remove contaminated clothing and wash clothing before reuse.

USER SAFETY REQUIREMENTS

- User should wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- Remove clothing immediately if pesticide soaks clothing. Change clothing as soon as possible after use.
- Wash the outside of gloves before removing.
- Do not allow children or pets to contact area until spray has dried.

First Aid

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact the National Pesticide Information Center at 1-800-858-7378 for emergency medical treatment information.

If in eyes: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.

If on skin or clothing: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.

If swallowed: Call a poison control center or doctor immediately for treatment advice. Do not give any liquid to the person. Do not induce vomiting unless told to do so by the poison control center or doctor. Do not give anything by mouth to an unconscious person.

If inhaled: Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control center or doctor for further treatment advice.

NOTE TO PHYSICIAN: Probable mucosal damage may contraindicate the use of gastric lavage. This product may pose an aspiration pneumonia hazard. Contains petroleum distillate. This product may cause cholinesterase inhibition. Atropine is antidotal.

Environmental Hazards

This pesticide is toxic to fish. Do not apply directly to water. Do not contaminate water when disposing of equipment washwaters or rinsate. This product is highly

toxic to bees exposed to direct treatment on blooming crops or weeds. Do not apply this product or allow it to drift to blooming crops or weeds if bees are visiting the treatment area.

Physical & Chemical Hazards

Extremely flammable. Keep away from fire, sparks and heated surfaces.

STORAGE & DISPOSAL

Storage: Store in a cool, dry area inaccessible to children or pets. Store and transport in an upright position.

Disposal: If empty: Do not reuse this container. Place in trash or offer for recycling if available. **If partly filled:** Call your local solid waste agency for disposal instructions. Never place unused product down any indoor or outdoor drain.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Not for use on plants being grown for sale or other commercial use, or for commercial seed production, or for research purposes.

GORDON'S LIQUID FRUIT TREE SPRAY is a complete concentrate containing insecticide, fungicide, aphicide, miticide, scabicide and spreader sticker. Easy to use; mixes with water instantly; no plugged nozzles; no messy powders to handle, measure or mix; no pre-mixing or straining necessary. Designed especially for home orchards to protect apples, cherries, grapes, peaches, strawberries, roses, evergreens and flowers from the ravages of insects and diseases listed.

SHAKE PRODUCT THOROUGHLY BEFORE USE. Contains micronic particles which settle upon standing and require reblending by agitation. Choose a cool, calm period, preferably early morning or evening, when trees will not be subjected to hot direct rays of the sun. Do not apply if rain is expected before leaf surfaces dry. It is easier to prevent disease and scale damage than to cure same, therefore, a preventative 10-14 day spray schedule is recommended. When insects are present, use maximum application rate. When active disease spores are present (warm humid days), shorten spray intervals to 7-10 days at maximum application rate. Apply spray to leaves, fruit branches, trunk and base of tree until lightly dripping. Shake sprayer occasionally or agitate to keep spray particles in suspension during application. Carefully read and follow the spray schedule.

Not for use on plants being grown for sale or other commercial use, or for commercial seed production, or for research purposes.

General Precautions & Restrictions

- **ENTRY RESTRICTIONS:** Do not allow people or pets to enter the treated area until sprays have dried.
- Do not apply this product in a way that will contact any person or pet, structures people occupy at any time and the associated property, parks and recreation areas, nontarget crops, aquatic and wetland areas, woodlands, pastures or rangelands, either directly or through drift. Keep people and pets out of the area during application.
- Apply when air is calm (not more than 10 mph) using largest spray droplet size possible to avoid drift and contact with eyes and skin.
- Do not apply to residential lawns.
- Do not apply this product through any type of irrigation equipment.

Specific Directions for use: Use 1 to 2 pts. spray suspension on 5 to 8 ft. trees. Use 1 to 2 qts. spray suspension on 8 to 14 ft. trees. On large trees use 1 to 3 gals. spray suspension.

Start spraying at the farthest corner of the treatment area and work backward to avoid contact with wet surfaces. Allow spray to dry in treated areas before reentering. For trees taller than 10 feet, consider hiring a licensed professional. Spray thoroughly to wet upper and lower leaf surfaces, stems and branches.

Do not repeat applications more than indicated in these directions for use.

APPLES

Insects/Diseases: Apple maggots, codling moths, Japanese beetles, leaf hoppers, plum curculios, aphids, bud moth, Forbes scale, eastern tent caterpillar, red banded leaf roller, mites, bitter rot, black rot, frog-eye leaf spot, Botryosphaeria (white rot), botrytis rot, bullseye rot, Brooks fruit spot, flyspeck, cedar rust, quince rust, scab, sooty blotch.

CHERRIES

Insects/Diseases: Plum curculios, cherry fruit flies, cherry fruit worms, black cherry aphid, fruit tree leaf roller, bud moth, Japanese beetle, Botrytis rot, brown rot blossom blight and fruit rot, cherry leaf spot.

GRAPES

Insects/Diseases: Flea beetles, Japanese beetles, grape leaf skeletonizers, rose chafers, grape berry moth, spider mites, mealybugs, leafhoppers, black rot, downy mildew (berries only), Botrytis rot.

PEACHES

Insects/Diseases: Plum curculios, Japanese beetle, spider mite, oriental fruit moth; green, peach, black cherry, black peach, rusty plum aphids; brown rot (fruit), rhizopus, scab, brown rot (blossom blight), cornynem blight.

FRUIT TREE STAGES

Dormant: Absence of growth fall, winter or spring.

Green Tip Stage: When branch tips show green tissue.

Pre Bloom Stage: When blossom buds show full color before opening.

Bloom: When blossoms are open. To protect pollinating insects, do not spray during blossoming period. Use another product that contains only a fungicide during the blossoming period, if disease control is needed.

Petal Fall Stage: When last petals have fallen from blossoms.

1st Cover: 7-10 days after petal fall.

2nd Cover to Harvest: To cut off date before harvest.

STRAWBERRIES

Insects/Diseases: Strawberry weevils, aphids, spider mites, lygus bugs, flea beetles, omnivorous leafhopper, spittle bugs, Japanese beetles, field crickets, Botrytis rot (gray mold), leaf spots. **Use 2½ teaspoons concentrate per gallon of water.** Allow 14 days between last application and harvest. Do not use more than 3 quarts of spray suspension per 100 sq.ft. of strawberries (broadcast spray) or per 100 linear feet of band or row 1 ft. wide (banded or directed) spray per single application. Make no more than 8 foliar applications per growing season.

ROSES, EVERGREENS AND FLOWERS

Mix 1½ Tablespoons per gallon of water.

Insects: Aphids, apple maggot, bagworm, black cutworm, bud moth, cherry fruit fly and worm, codling moth, plum curculios, flea beetles, fruit tree leaf roller, gypsy moth, Japanese beetle, leaf hoppers, lesser peach tree borer, mealybugs,

mites (clover, European red, two-spotted and Wilmette), oriental fruit moth, pear slugs, psylla, red banded leaf roller, scale (Forbes, Putman, San Jose), spittle bug, tent caterpillars, unspotted tentiform leaf miner, and yellow neck caterpillar.

Diseases: Specifically — CARNATION (Rust, Alternaria Leaf Spot), CHRYSANTHEMUMS (Botrytis Flower Blight), ROSE (Black Spot) — Apply at first sign of disease. Repeat at 7 to 10 day intervals. Use the shorter interval under conditions favorable to infection.

Begin applications when pests or disease symptoms first appear or conditions favor their development and repeat at weekly intervals or as necessary to maintain control. Remember, it is easier to prevent damage than to cure it. Therefore, a preventative spray schedule is recommended. Do not use if rain is expected shortly after application. Select still periods for application (early morning or late evening) to reduce waste by blow away and blow back applicator.

2 Tablespoons = 1 Fluid Ounce = 1/128 Gallon



795/10-2008 AP082405
EPA REG. NO. 4-122-33955

DISTRIBUTED BY
PBI/GORDON CORPORATION
1217 WEST 12TH STREET
KANSAS CITY, MISSOURI 64101
www.pbigordon.com

All rates in chart below are given in Tablespoons per gallon. Shake well before using.

SPRAY CHART						
Start and repeat at 10-14 day intervals up to 14 days before harvest.						
Fruit	Green Tip	Pre Bloom	Full Pink	Petal Fall	1st Cover	2nd Cover
Apple (1)	1½ tbs.	2½ tbs.	2½ tbs.	2½ tbs.	2½ tbs.	2½ tbs.
Cherry (2)	1½ tbs.	2½ tbs.	2½ tbs.	2½ tbs.	2½ tbs.	2½ tbs.
Grape (3)	1½ tbs.	2½ tbs.	2½ tbs.	2½ tbs.	2½ tbs.	2½ tbs.
Peach (1)	1½ tbs.	1½ tbs.	2½ tbs.	Repeat at 10-14 day intervals up to 21 days before harvest. 2½ tbs. 2½ tbs. 2½ tbs.		

- (1): Make no more than 8 foliar applications per growing season.
 (2): Make no more than 7 foliar applications per growing season.
 (3): Do not use more than 2 quarts of spray suspension per 100 sq. ft. of grape vine.
 Make no more than 7 foliar applications per growing season.

