

[CLICK HERE TO JUMP TO USE DIRECTIONS](#)



LIQUID WEED & FEED 15-0-0

Control Broadleaf Weeds and Fertilize in One Step!

ACTIVE INGREDIENTS:

Dimethylamine salt of 2,4-dichlorophenoxyacetic acid 4.51%
 Dimethylamine salt of (+)-R-2-(2-methyl-4-chlorophenoxy)propionic acid 2.33%

INERT INGREDIENTS: 93.16%
 TOTAL 100.00%

THIS PRODUCT CONTAINS:

0.35 lb. 2,4-dichlorophenoxyacetic acid equivalent per gallon or 3.75%.
 0.18 lb. (+)-R-2-(2-methyl-4-chlorophenoxy)propionic acid equivalent per gallon or 1.93%.

Contains the single isomer form of Mecoprop-p.
 Isomer Specific by AOAC Methods.

GUARANTEED ANALYSIS FOR 15-0-0

Total Nitrogen (N) 15%
 15% Urea Nitrogen
 Source: Derived From Urea.

KEEP OUT OF REACH OF CHILDREN CAUTION

See below for additional Precautionary Statements and First Aid.



READ THE ENTIRE LABEL FIRST. OBSERVE ALL PRECAUTIONS AND FOLLOW DIRECTIONS CAREFULLY.

PRECAUTIONARY STATEMENTS

Hazards to Humans & Domestic Animals

CAUTION: Harmful if swallowed. Avoid contact with eyes, skin or clothing. Wear goggles when applying this product. When using this product, wear long-sleeved shirt, long pants, socks, shoes and rubber gloves.

After using this product, rinse gloves before removing, remove clothing and launder separately before reuse, and promptly and thoroughly wash hands and exposed skin with soap and water. Remove saturated clothing as soon as possible and shower.

Avoid contamination of food or feedstuffs.

First Aid

If in eyes:	<ul style="list-style-type: none"> • Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing. • Call a poison control center or doctor for treatment advice.
If on skin or clothing:	<ul style="list-style-type: none"> • Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15-20 minutes. • Call a poison control center or doctor for treatment advice. <p style="text-align: right;"><i>(cont. on next column)</i></p>

First Aid (continued)

If swallowed:	<ul style="list-style-type: none"> • Call a poison control center or doctor immediately for treatment advice. • Have person sip a glass of water if able to swallow. • Do not induce vomiting unless told to do so by a poison control center or doctor. • Do not give anything by mouth to an unconscious person.
Have the product container or label with you when calling a poison control center or doctor or going for treatment. You may also call toll-free, 1-877-800-5556 for emergency medical treatment information.	

ENVIRONMENTAL HAZARDS:

This product is toxic to aquatic invertebrates. Drift or runoff may adversely affect aquatic invertebrates and nontarget plants. Do not apply directly to water. Do not contaminate water when disposing of equipment washwater. Do not apply when weather conditions favor drift from target area.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Do not allow people (other than applicator) or pets on treatment area during application. Do not enter treatment area until spray has dried or dust has settled.

USE PRECAUTIONS:

Do not allow spray to drift to vegetables, flowers, ornamental plants, shrubs, trees and other desirable plants since injury may result. Do not spray when grass is emerging from dormancy. Do not apply to newly seeded lawn grasses until well established. Do not spray when the temperature exceeds 85°F. Seed can be sown 3 to 4 weeks after application. Do not use on carpetgrass, dichondra, St. Augustinegrass, centipede grass, closely-mowed bentgrass, nor on lawns or turf where desirable clovers are present. Do not spray roots of trees and ornamentals. Do not exceed specified dosages for any area. Be particularly careful within dripline of tree and other ornamental species.

- WEEDS -

- | | |
|------------------|-----------------------|
| Black medic | Mallow |
| Dock | Mouseear chickweed |
| Clover | Oxalis |
| Common chickweed | Pigweed |
| Dandelion | Plantain |
| Dock | Purslane |
| English Daisy | Ragweed |
| Knotweed | and many other |
| Lambsquarters | broadleaf lawn weeds. |

A lawn will typically utilize 1 to 4 pounds of nitrogen per 1,000 square feet each growing season. When applied as directed this product supplies 0.07 pounds of nitrogen per 1,000 square feet with each application. Use this product in conjunction with an additional fertilization program to provide at least 1 pound of nitrogen per 1,000 square feet each growing season. Consult your state agricultural extension agent for the proper amount of nitrogen to be applied in your area.

WHERE TO USE:

Gordon's Liquid Weed & Feed may be used on bentgrass, bermudagrass, bluegrass, fescue lawns, vacant lots, playgrounds and other similar areas where broadleaf weeds are present. Gordon's Liquid Weed & Feed controls most broadleaf weeds, including chickweed, clover, dandelion, knotweed and plantain.

WHEN TO USE:

1. Spray at any time during the growing season when weeds are actively growing.
2. On new lawns, wait until the grass has reached a height of 2 inches before making an overall broadcast spray treatment. Wait 3 weeks after application before seeding or planting stolons.
3. Poor weed control may result if spray is applied during drought or just before rain.
4. For best results, mow 1 to 2 days before treating.
5. Do not water lawn within 24 hours after treatment.

The maximum application rate to turf is 0.8 pounds 2,4-D acid equivalent per acre per application per site. The maximum number of broadcast applications per treatment site is 2 per year.

SHAKE WELL BEFORE USING

MIXING INSTRUCTIONS WITH WATER (for pull-behind sprayers)

Mix one quart of Gordon's Liquid Weed and Feed with 5 gallons of water and apply over 5,000 square feet.

For larger areas:

- Use 3 quarts in 15 gallons of water and apply to 15,000 square feet.
- Use 5 quarts in 25 gallons of water and apply to 25,000 square feet.
- Use 6 quarts in 30 gallons of water and apply to 30,000 square feet.

PESTICIDE STORAGE: Keep from freezing. Store in original container in a locked storage area inaccessible to children. Do not store near food or feedstuffs.

CONTAINER DISPOSAL:

If Empty — Do not reuse this container. Place in trash or offer for recycling if available.

If Partly Filled — Call your local solid waste agency for disposal instructions. Never place unused product down any indoor or outdoor drain.

LIMITED WARRANTY AND DISCLAIMER

The manufacturer warrants only that the chemical composition of this product conforms to the ingredient statement given on the label, and that the product is reasonably suited for the labeled use when applied according to the Directions for Use.

THE MANUFACTURER NEITHER MAKES NOR INTENDS ANY OTHER EXPRESS OR IMPLIED WARRANTIES, INCLUDING ANY WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE, WHICH ARE EXPRESSLY DISCLAIMED. This limited warranty does not extend to the use of the product inconsistent with label instructions, warnings or cautions, or to use of the product under abnormal conditions such as drought, excessive rainfall, tornadoes, hurricanes, etc. These factors are beyond the control of the manufacturer or the seller. Any damages arising from a breach of the manufacturer's warranty shall be limited to direct damages, and shall not include indirect or consequential damages such as loss of profits or values, except as otherwise provided by law.

The terms of this Limited Warranty and Disclaimer cannot be varied by any written or verbal statements or agreements. No employee or agent of the seller is authorized to vary or exceed the terms of this Limited Warranty and Disclaimer in any manner.

Information regarding the contents and levels of metals in this product is available on the internet at <http://www.aapfco.org/metals>

GORDON'S is a registered trademark of PBI/Gordon Corporation. FL #987

714/1-2006 AP112205
EPA REG. NO. 2217-742

Manufactured By

pbi/gordon
CORPORATION
An Employee-Owned Company
1217 West 12th Street
Kansas City, Missouri 64101
www.pbigordon.com