

CLICK HERE TO JUMP TO USE DIRECTIONS



TRIMEC® 959 BROADLEAF HERBICIDE

For use by professional turf maintenance personnel, landscaping or commercial applicators only.

Controls Dandelion, Chickweed, Knotweed, Plantain, Henbit, Spurge and many other broadleaf weeds.



ACTIVE INGREDIENTS:

Dimethylamine salt of 2-methyl-4-chlorophenoxyacetic acid . . .	38.68%
Dimethylamine salt of (+)-(R)-2-(2-methyl-4-chlorophenoxy) propionic acid	8.16%
Dimethylamine salt of dicamba: 3,6-dichloro-o-anisic acid	3.81%
INERT INGREDIENTS:	<u>49.35%</u>
TOTAL 100.00%	

THIS PRODUCT CONTAINS:

- 2.97 lbs. 2-methyl-4-chlorophenoxyacetic acid equivalent per gallon or 31.59%.
- 0.63 lbs. (+)-(R)-2-(2-methyl-4-chlorophenoxy) propionic acid equivalent per gallon or 6.74%.
- 0.29 lbs. 3,6-dichloro-o-anisic acid equivalent per gallon or 3.16%.

Contains single isomer form of Mecoprop-p.

TRIMEC® is a registered trademark of PBI/GORDON CORPORATION.

**KEEP OUT OF REACH OF CHILDREN
DANGER**

See below for additional Precautionary Statements and First Aid.

KEEP FROM FREEZING



READ THE ENTIRE LABEL FIRST. OBSERVE ALL PRECAUTIONS AND FOLLOW DIRECTIONS CAREFULLY.

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS & DOMESTIC ANIMALS

DANGER: Corrosive. This concentrate causes irreversible eye damage. Do not get in eyes, on skin, or on clothing. Avoid exposure to spray mist. Harmful or fatal if swallowed or absorbed through skin. Harmful if inhaled. Wear long pants, long-sleeved shirt, shoes, gloves, and face shield or goggles or shielded safety glasses when mixing, or loading this concentrated product. It is recommended that safety glasses include front, brow and temple protection. After product has been diluted in accordance with use directions below, protective eyewear and long-sleeved shirt are not necessary.

Wash thoroughly with soap and water after handling and before eating or smoking. Remove contaminated clothing and wash before reuse.

FIRST AID

If in eyes:	<ul style="list-style-type: none"> • Hold eye open and rinse slowly and gently with water for 15-20 minutes. • Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. • Call a poison control center or doctor for treatment advice. <i>(Continued on next column)</i>
--------------------	---

FIRST AID (continued)

If on skin:	<ul style="list-style-type: none"> • Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15-20 minutes. • Call a poison control center or doctor for treatment advice.
If swallowed:	<ul style="list-style-type: none"> • Call a poison control center or doctor immediately for treatment advice. • Have person sip a glass of water if able to swallow. • Do not induce vomiting unless told to do so by the poison control center or doctor. • Do not give anything by mouth to an unconscious person.
If inhaled:	<ul style="list-style-type: none"> • Move person to fresh air. • If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible. • Call a poison control center or doctor for treatment advice.
<p>Have the product container or label with you when calling a poison control center or doctor or going for treatment. You may also contact 1-877-800-5556 for emergency medical treatment advice.</p> <p>NOTE TO PHYSICIAN: Probable mucosal damage may contraindicate the use of gastric lavage.</p>	

ENVIRONMENTAL HAZARDS: Drift or runoff may adversely affect non-target plants. Do not apply directly to water, or to areas where surface water is present, or to intertidal areas below mean high water mark. Do not contaminate water when disposing of equipment washwater or rinsate. When cleaning equipment, do not pour the washwater on the ground; spray or drain over a large area away from wells and other water sources. Do not apply when weather conditions favor drift from target area.

Most cases of groundwater contamination involving phenoxy herbicides such as MCPP-p and MCPA have been associated with mixing/loading and disposal sites. Caution should be exercised when handling MCPP-p and MCPA pesticides at such sites to prevent contamination of groundwater supplies. Use of closed systems for mixing and transferring this pesticide will reduce the probability of spills. Placement of the mixing/loading equipment on an impervious pad to contain spills will help prevent groundwater contamination.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Do not reenter or permit workers to reenter treated area until sprays have dried or dusts have settled.

Not for use on turf being grown for sale or other commercial use as sod, or for commercial seed production, or for research purposes.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

STORAGE: Store in original container in a locked storage area inaccessible to children or pets. Keep from freezing. To prevent cross-contamination, do not store near herbicides, fertilizers, insecticides, fungicides, or seeds.

PESTICIDE DISPOSAL: Pesticide wastes are acutely hazardous. Improper disposal of excess pesticide, spray mixture or rinsate is a violation of Federal law and may contaminate groundwater. If these wastes cannot be disposed of by use according to label instructions, contact your state Pesticide or Environmental Control Agency or the Hazardous Waste Representative at the nearest EPA Regional Office for guidance.

CONTAINER DISPOSAL: For Plastic Containers: Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke. **For Metal Containers:** Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by other procedures approved by state and local authorities.

GENERAL INFORMATION

Trimec® 959 Broadleaf Herbicide provides selective broadleaf control in warm season and cool season turfgrass. Established turf of warm season grasses include, but are not limited to, bermudagrass, bahia-

grass, and zoysiagrass. Established turf of cool season grasses include, but are not limited to, Kentucky bluegrass, perennial ryegrass, tall fescue, and turf-type tall fescues. Do not apply this product on carpetgrass, dichondra, or established turf containing desirable clovers.

USE PRECAUTIONS

Avoid drift of spray mist onto vegetables, flowers, ornamental plants, shrubs, trees and other desirable plants. Do not pour spray solutions near these plants. Do not spray on carpetgrass, dichondra nor on lawns or turf where desirable clovers are present. Use only lawn type sprayers. NOTE: Coarse sprays are less likely to drift. Avoid application to exposed active feeder roots of shallow-rooted trees such as poplar species and sensitive ornamentals. Do not exceed specified dosages for any area. Be particularly careful within the dripline of trees and other ornamental species. Do not spray bermudagrass or zoysiagrass while emerging from dormancy, unless injury can be tolerated. Applications to newly seeded lawns may cause injury. Delay application until well established or after two or three mowings. Do not use broadcast applications when air temperatures exceed 85°F; some injury may be expected with spot treatments when air temperatures exceed 85°F. Seed can be sown 3 to 4 weeks after application. Do not contaminate domestic or irrigation waters. Some injury can be expected to turf when used on centipede-grass or St. Augustinegrass; only spot spray weeds on these types of grasses. Clover will be damaged if sprayed. After using this product, clean sprayer with soap or detergent and water, and rinse thoroughly before applying other pesticides. Failure to observe the above precautions may result in injury to desirable plants. Do not apply this product through any type of irrigation system.

ORNAMENTAL LAWNS & TURF (other than bentgrass): Apply 3 to 4¼ pints of product in 20 to 240 gallons of water per acre. Maximum control of weeds will be obtained from spring or early fall applications when weeds are actively growing. Avoid applying during long excessively dry or hot periods unless adequate irrigation is available. Do not apply to newly seeded turf until well established or after the second or third mowing. Grass seed can be sown 3 to 4 weeks after application at the recommended rates.

BENTGRASS TURF: In closely-mowed bentgrass (putting and bowling greens) apply at a maximum rate of 1 fluid ounce of product per 1,500 square feet, preferably in May or mid-August through September. Slight turf yellowing will disappear after about one week.

NOTE: Care should be taken to avoid overdosing bentgrass or injury may result. High spray volumes (i.e. 1 fluid ounce of product in 5 gallons of water per 1,500 square feet) will aid in obtaining uniform coverage. If hand operated sprayers are used, it is preferable to use a single nozzle sprayer rather than a multiple nozzle boom. Sweeping movements with a multiple nozzle boom will result in heavy local over-application and subsequent turf discoloration or injury.

BROADLEAF WEEDS

Aster, white heath & white prairie	Field bindweed (*morningglory & creeping jenny)
Bedstraw	Field oxeye-daisy
Beggarweed, creeping	(*creeping oxeye)
Bindweed	Filaree, whitestem & redstem
Black medic	Florida pusley
Broadleaf plantain	Ground ivy
Buckhorn plantain	Groundsel
Bull thistle	Hawkweed
Burclover	Healall
Burdock, common	Henbit
Buttercup, creeping	Lambsquarters
Carpetweed	Lawn burweed
Chickweed, common	Lespedeza, common
Chicory	Mallow, common
Cinquefoil	Matchweed
Clover	Mouseear chickweed
Compassplant	Oxalis (*yellow woodsorrel & creeping woodsorrel)
Curly dock	Parsley-piert
Dandelion	Pennsylvania smartweed
Dayflower	Pennywort (*dollarweed)
Deadnettle	Pepperweed
Dock	Pigweed
Dogfennel	Pineappleweed
English daisy	Plantain
False dandelion	Poison ivy
(*spotted catsear & common catsear)	Poison oak

Prostrate knotweed (*knotweed)	White clover (*Dutch clover, honeysuckle clover, white trefoil & purplewort)
Puncturevine	Wild carrot
Purslane	Wild garlic
Ragweed	Wild geranium
Red sorrel (*sheep sorrel)	Wild mustard
Shepherdspurse	Wild onion
Spotted spurge	Wild strawberry
Spurge	Wild violet
Sunflower	Yarrow
Thistle	Yellow rocket
Velvetleaf (*pie marker, Indian mallow)	
Veronica (*corn speedwell)	
Virginia buttonweed	*Synonyms

LIMITED WARRANTY AND DISCLAIMER

The manufacturer warrants only that the chemical composition of this product conforms to the ingredient statement given on the label, and that the product is reasonably suited for the labeled use when applied according to the Directions for Use.

THE MANUFACTURER NEITHER MAKES NOR INTENDS ANY OTHER EXPRESS OR IMPLIED WARRANTIES, INCLUDING ANY WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE, WHICH ARE EXPRESSLY DISCLAIMED.

This limited warranty does not extend to the use of the product inconsistent with label instructions, warnings or cautions, or to use of the product under abnormal conditions such as drought, excessive rainfall, tornadoes, hurricanes, etc. These factors are beyond the control of the manufacturer or the seller. Any damages arising from a breach of the manufacturer's warranty shall be limited to direct damages, and shall not include indirect or consequential damages such as loss of profits or values, except as otherwise provided by law.

The terms of this Limited Warranty and Disclaimer cannot be varied by any written or verbal statements or agreements. No employee or agent of the seller is authorized to vary or exceed the terms of this Limited Warranty and Disclaimer in any manner.

Manufactured By



An Employee-Owned Company

1217 West 12th Street
Kansas City, Missouri 64101

www.pbigordon.com