

[CLICK HERE TO JUMP TO USE DIRECTIONS](#)

GORDON'S® TRIMEC® CRABGRASS PLUS LAWN WEED KILLER READY SPRAY

*Also Kills Dandelion & Clover
Plus 200 Other Listed Weeds*

ACTIVE INGREDIENTS:	
2,4-D, dimethylamine salt	6.42%
Quinclorac	2.13%
Dicamba, dimethylamine salt	0.60%
INERT INGREDIENTS:	90.85%
TOTAL 100.00%	

THIS PRODUCT CONTAINS:
 0.456 lb. 2,4-dichlorophenoxyacetic acid equivalent per gallon or 5.33%
 0.182 lb. 3,7-dichloro-8-quinolinecarboxylic acid per gallon or 2.13%
 0.043 lb. 3,6-dichloro-o-anisic acid equivalent per gallon or 0.50%
 Isomer Specific By AOAC Methods.

TRIMEC® is a registered trademark of PBI/Gordon Corporation.

KEEP OUT OF REACH OF CHILDREN CAUTION



**READ THE ENTIRE LABEL FIRST.
OBSERVE ALL PRECAUTIONS AND
FOLLOW DIRECTIONS CAREFULLY.**

PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals

CAUTION: Causes moderate eye irritation. Avoid contact with eyes, skin or clothing. Harmful if swallowed. When using this product, wear long-sleeved shirt, long pants, socks, shoes, and rubber gloves.

Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum or using tobacco. After using this product, rinse gloves before removing, remove clothing and launder separately before reuse, and promptly and thoroughly wash hands and exposed skin with soap and water. Remove saturated clothing as soon as possible and shower.

User Safety Recommendations

- Users should wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Users should remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing. If pesticide gets on skin, wash immediately with soap and water.
- Users should remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

First Aid

If in eyes:	<ul style="list-style-type: none"> • Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first minutes, then continue rinsing eye. • Call a poison control center or doctor for treatment advice.
If swallowed:	<ul style="list-style-type: none"> • Call a poison control center or doctor immediately for treatment advice. • Have person sip a glass of water if able to swallow. • Do not induce vomiting unless told to by a poison control center or doctor. • Do not give anything by mouth to an unconscious person.

Have the product container or label with you when calling a poison control center or doctor or going for treatment. You may also contact 1-877-800-5556 for emergency medical treatment advice.

Environmental Hazards

This pesticide is toxic to fish and aquatic invertebrates. To protect the environment, do not allow pesticide to enter or run off into storm drains, drainage ditches, gutters or surface waters. Applying this product in calm weather when rain is not predicted for the next 24 hours will help to ensure that wind or rain does not blow or wash pesticide off the treatment area.

This chemical has properties and characteristics associated with chemicals detected in groundwater. The use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination. Application around a cistern or well may result in contamination of drinking water or groundwater.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

GENERAL PRECAUTIONS AND RESTRICTIONS:

- Do not apply this product in a way that will contact any person or pet, either directly or through drift. Keep people and pets out of the area during application.
- Do not allow people or pets to enter the treated area until sprays have dried.

SHAKE WELL BEFORE USING

1. WHAT IT DOES: Kills major broadleaf and troublesome grass weeds including dandelion, chickweed and clover as well as crabgrass and foxtail. (See below for complete list.)

2. HOW IT WORKS: Trimec® Crabgrass Plus Lawn Weed Killer Ready Spray contains three proven weed killers that target lawn weeds and crabgrass. This herbicide enters the lawn weeds through their leaves and moves throughout the plant to provide control. And lawns listed on this label will not be harmed when used as directed.

3. WHERE TO USE: Trimec Crabgrass Plus Lawn Weed Killer Ready Spray can be applied to residential lawns with the following turfgrass species:

Lawns To Be Treated:	
Cool-Season Turfgrass	Warm-Season Turfgrass
Kentucky bluegrass, Ryegrass (perennial and annual), Tall fescue, Mixtures of cool-season grasses containing fine fescues (red, hard and chewings)	Bermudagrass ¹ (common and hybrid), Zoysiagrass, Buffalograss
¹ Application to bermudagrass may cause temporary yellowing or discoloration but full recovery can be expected. Do not apply to bermudagrass when daytime temperatures exceed 85°F.	

DO NOT apply Trimec Crabgrass Plus Lawn Weed Killer Ready Spray to the following:

Bahiagrass, Bentgrass (colonial and creeping), Carpetgrass, Centipedegrass, Dichondra, Gardens including vegetables, fruit trees, vines and berries, Ornamental plants (flowers, trees, groundcovers, landscape beds and shrubs), Seashore paspalum, St. Augustinegrass, Turfgrass species that are not listed on this label.

4. GRASS AND BROADLEAF WEEDS CONTROLLED:

A partial list is presented below:

GRASS WEEDS		
Crabgrass, large and smooth	Foxtail, green, yellow and giant	Signalgrass, broadleaf
BROADLEAF WEEDS		
Annual yellow sweetclover	Bull thistle	Cudweed
Aster	Bullnettle	Curly dock
Austrian fieldcress	Burclover	Daisy, English
Bedstraw	Burdock	Daisy fleabane
Beggarticks	Burning nettle	Daisy, oxeye
Betony, Florida	Burweed	Dandelion
Bindweed, field	Buttercup	Dichondra
Bird vetch	Buttonweed	Dogbane
Bitter wintercress	Canada thistle	Dogfennel
Bittercress, hairy	Carolina geranium	Dollarweed
Bitterweed	Carpetweed	Elderberry
Black-eyed Susan	Catchweed bedstraw	False dandelion
Black medic	Catnip	Falseflax
Black mustard	Catsear	False sunflower
Blackseed plantain	Chickweed, common	Fiddleneck
Blessed thistle	Chickweed, mouseear	Florida pusley
Bloodflower milkweed	Chicory	Frenchweed
Blue lettuce	Cinquefoil	Galinsoga
Blue vervain	Clover, crimson	Goathead
Bracted plantain	Clover, hop	Goldenrod
Brassbuttons	Clover, red	Ground ivy
Bristly oxtongue	Clover, strawberry	Gumweed
Broadleaf dock	Clover, sweet	Hairy fleabane
Broadleaf plantain	Clover, white	Hawkweed
Broomweed	Cockle	Healall
Buckhorn	Cocklebur	Heartleaf drymary
Buckhorn plantain	Common mullein	Heathaster
Bulbous buttercup	Creeping Jenny	Hedge bindweed
	(Creeping Charlie)	

(continued on next page)

BROADLEAF WEEDS (cont.)

Hedge mustard	Poorjoe	Thistle
Hemp	Prairie sunflower	Trailing crownvetch
Henbit	Prickly lettuce	Tumble mustard
Hoary cress	Prickly sida	Tumble pigweed
Hoary plantain	Prostrate knotweed	Velvetleaf
Hoary vervain	Prostrate pigweed	Venice mallow
Horsenettle	Prostrate spurge	Virginia buttonweed
Jimsonweed	Prostrate vervain	Virginia creeper
Knawel	Puncturevine	Virginia pepperweed
Knotweed	Purslane, common	Water pennywort
Kochia	Ragweed	Wavyleaf bullthistle
Lambsquarters	Red sorrel	Western clematis
Lespedeza	Redroot pigweed	Western salsify
Mallow	Redstem filaree	White mustard
Matchweed	Rough cinquefoil	Wild aster
Mexicanweed	Rough fleabane	Wild buckwheat
Milk vetch	Russian pigweed	Wild carrot
Morningglory	Russian thistle	Wild four-o'clock
Mouseear hawkweed	Scarlet pimpernel	Wild garlic
Mugwort	Scotch thistle	Wild geranium
Musk thistle	Sheep sorrel	Wild lettuce
Mustard	Shepherdspurge	Wild marigold
Narrowleaf plantain	Slender plantain	Wild mustard
Narrowleaf vetch	Smallflower galinsoga	Wild onion
Nettle	Smooth dock	Wild parsnip
Orange hawkweed	Smooth pigweed	Wild radish
Oriental cocklebur	Sorrel	Wild rape
Oxalis	Sowthistle	Wild strawberry
Parsley-piert	Spanishneedles	Wild sweet potato
Parsnip	Speedwell	Wild vetch
Pearlwort	Spiny amaranth	Wild violet
Pennycress	Spiny cocklebur	Woodsorrel
Pennywort	Spiny sowthistle	Woolly croton
Peppergrass	Spotted catsear	Woolly morningglory
Pepperweed	Spotted spurge	Woolly plantain
Pigweed	Spurweed	Wormseed
Pineywoods bedstraw	Stinging nettle	Yarrow
Plains coreopsis	Strawberry, India mock	Yellow rocket
(tickseed)	Tall nettle	Yellowflower
Plantain	Tall vervain	pepperweed
Poison ivy	Tansy ragwort	and other
Poison oak	Tansy mustard	broadleaf weeds
Pokeweed	Tanweed	

5. WHEN TO USE:

Apply this product when weeds are small and actively growing.

Weed species	Season	Timing	Maximum number of applications ¹	Application interval, days	Comments
Crabgrass (large and smooth):	Spring	From crabgrass emergence until the crabgrass is 3 to 4 inches in height.	2	14	Re-treat at the same rate at 14 days later for dense populations or for continued crabgrass emergence.
Foxtail (green, giant and yellow):	Spring and early summer	During the early growth stages or when foxtail is 1 to 4 inches in height.	2	14	Re-treat at the same rate at 14 days later for dense populations.
Broadleaf signalgrass	Spring and early summer	During the early growth stages or during the 3 to 7 leaf stage.	2	14	Re-treat at the same rate at 14 days later if needed.
Dandelion, plantain, buttercup, speedwell, knotweed and other listed broadleaf weeds	Spring	When broadleaf weeds are small and actively growing.	2	—	Apply when broadleaf weeds are young and actively growing with adequate soil moisture.
Henbit, chickweed, white clover, ground ivy, lawn burweed and other listed broadleaf weeds	Fall	When broadleaf weeds are young and actively growing.	2	—	Apply when broadleaf weeds are young and actively growing with adequate soil moisture.

¹The maximum application rate is 6.4 fl. oz. of product per 1,000 sq. ft. per application (1.0 lb. 2,4-D acid equivalent per acre per application).

The maximum number of broadcast applications is limited to 2 per year. Do not apply more than 2 applications per year. The maximum seasonal rate is 12.8 fl. oz. of product per 1,000 sq. ft. (2.0 lbs. 2,4-D acid equivalent per acre).

6. HOW MUCH TO USE:

This retail sprayer bottle will cover 5,000 sq. ft. when this product is applied at 6.4 fl. oz. of product per 1,000 sq. ft. Over-application or rates above those recommended on this label can cause turf injury.

Instructions:

- Determine (Measure) the total lawn area to be sprayed. For rectangular areas, multiply the length by the width to obtain total square feet.
- **CONNECT:** Remove bottle cap and screw the spray nozzle onto the bottle. Attach garden hose to spray nozzle. Turn the spray nozzle lever to the "OFF" position.
- **SPRAY:** Turn on water at faucet. Extend hose to the farthest area of lawn to be treated. To begin spraying, turn the spray nozzle lever to the "SPRAY" position. The herbicide mixes automatically with the water. Walk at a steady pace while spraying evenly and working towards the faucet.
- **FINISH:** To stop spraying, turn spray nozzle lever to the "OFF" position. Turn off water at faucet. To relieve pressure, turn spray nozzle to the "ON" position and point sprayer away from any desirable plants before disconnecting the sprayer nozzle from the garden hose.

7. CULTURAL TIPS AND NOTES:

When to Apply:	For established lawns For newly seeded areas For newly sodded, sprigged, or plugged areas	<ul style="list-style-type: none"> • Apply when daytime temperatures are below 90°F. • Apply this product to newly seeded grasses when well-established (approximately 6 weeks after seedling emergence) or after the third mowing. • Apply this product 3 to 4 weeks after the sodding, sprigging, or plugging operations.
Seeding:		<ul style="list-style-type: none"> • The treated area can be seeded at 4 weeks after the application of this product.
Irrigation:	Non-stressed turfgrass Stress from high temperatures, drought, insects or diseases	<ul style="list-style-type: none"> • Adequate soil moisture and favorable growing conditions enhance the performance of this product. • Avoid applications of this product when turfgrasses are under stress since injury may result. If dry conditions exist, schedule irrigations (watering) before and after the application. For best results, delay the irrigation until approximately 24 hours after application of this product.
Mowing:		<ul style="list-style-type: none"> • Delay mowing 2 days before and until 2 days after the application of this product. • Do not use clippings from the treated areas as mulch or compost around flowers, ornamentals, trees, or in vegetable gardens.
Nontarget plants:		<ul style="list-style-type: none"> • Do not apply this product to food crops (gardens, fruits and vegetables), forage crops, or ornamental plants including flowers, trees, shrubs, hedges, woody ornamentals, groundcovers established in landscape plantings and other nontarget plants. • Do not spray exposed roots of ornamentals and trees.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

PESTICIDE STORAGE: Keep from freezing. Store in original container in a locked storage area inaccessible to children or pets. Avoid contamination of food or feedstuffs.

PESTICIDE DISPOSAL AND CONTAINER HANDLING: Nonrefillable container. Do not reuse or refill this container. **If empty:** Place in trash or offer for recycling if available. **If partly filled:** Call you local solid waste agency for disposal instructions. Never place unused product down any indoor or outdoor drain.

LIMITED WARRANTY AND DISCLAIMER

FOR USE ONLY AS DIRECTED. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THE MANUFACTURER NEITHER MAKES NOR INTENDS ANY EXPRESS OR IMPLIED WARRANTIES, INCLUDING ANY WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE, WHICH ARE HEREBY EXPRESSLY DISCLAIMED. IN NO CASE SHALL THE MANUFACTURER BE LIABLE FOR INCIDENTAL, CONSEQUENTIAL, OR SPECIAL DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT. If these terms are not acceptable, return this product unopened immediately to the point of purchase, and the purchase price will be refunded in full. The terms of this Limited Warranty and Disclaimer cannot be varied by any written or verbal statements or agreements at the point of sale or elsewhere.

761/8-2009 AP081209
EPA REG. NO. 2217-896



MANUFACTURED BY
PBI/GORDON CORPORATION
1217 WEST 12TH STREET
KANSAS CITY, MISSOURI 64101
www.pbigordon.com