

CLICK HERE TO JUMP TO USE DIRECTIONS

GORDON'S®

TRIMEC® READY SPRAY LAWN WEED KILLER

**JUST CONNECT TO GARDEN HOSE
AND SPRAY! COVERS UP TO
5,000 SQ. FT. OF NORTHERN GRASSES.**

ACTIVE INGREDIENTS:

2,4-D, dimethylamine salt 5.56%
Mecoprop-p, dimethylamine salt 1.34%
Dicamba, dimethylamine salt 0.62%

INERT INGREDIENTS: 92.48%

THIS PRODUCT CONTAINS: TOTAL 100.00%

4.62% 2,4-dichlorophenoxyacetic acid equivalent.
1.11% (+)-(R)-2-(2-methyl-4-chlorophenoxy) propionic acid equivalent.
0.51% 3,6-dichloro-o-anisic acid equivalent.
Isomer Specific by AOAC Method.

Contains the single isomer form of Mecoprop-p.
TRIMEC® is a registered trademark of PBI/GORDON CORPORATION.
2,4-D dimethylamine salt (CAS Registry No. 2008-39-1)
Mecoprop-p, dimethylamine salt (CAS Registry No. 66423-09-4)
Dicamba, dimethylamine salt (CAS Registry No. 2300-66-5)

KEEP OUT OF REACH OF CHILDREN CAUTION

See below for additional *Precautionary Statements and First Aid.*
Shake well before using.



**READ THE ENTIRE LABEL FIRST.
OBSERVE ALL PRECAUTIONS AND
FOLLOW DIRECTIONS CAREFULLY.**

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Do not allow people (other than applicator) or pets on treatment area during application. Do not enter treatment area until spray has dried or dust has settled.

STORAGE: Keep from freezing. Store in original container in a locked storage area inaccessible to children and pets. Do not store near food or feedstuffs.

DISPOSAL:

If Empty – Do not reuse this container. Place in trash or offer for recycling if available.

If Partly Filled – Call your local solid waste agency for disposal instructions. Never place unused product down any indoor or outdoor drain.

This product is designed for use through this sprayer for controlling of broadleaf weeds in bahiagrass, bermudagrass, Kentucky bluegrass, fescue, perennial ryegrass, zoysiagrass, centipedegrass, and St. Augustinegrass lawns, vacant lots, playgrounds and other similar areas.

Temporary injury may occur on bentgrass, bermudagrass, and red fescue lawn grasses. Do not apply to carpetgrass, dichondra, Lippia, and lawns where desirable clovers are present.

Water lawn thoroughly 1 to 2 days before application.

Spray when air is calm to avoid spray drift that might injure desirable ornamental plants. Adjust water pressure at faucet to give coarse spray which is less likely to drift. Do not apply when temperatures are expected to exceed 85°F within 24 hours following application.

Newly seeded lawns may be treated after they have been mowed at least three times.

WHERE TO USE:

Trimec® Ready Spray Lawn Weed Killer may be used on residential and ornamental lawns, vacant lots, playgrounds and other similar areas where broadleaf weeds are present. Established turf of bahiagrass, bermudagrass, Kentucky bluegrass, centipedegrass, fescues, perennial ryegrass, St. Augustinegrass, and zoysiagrass may be treated. This product is especially effective on all hard-to-kill weeds such as poison ivy, poison oak, sheep sorrel and spurge. In addition, Trimec® Ready Spray Lawn Weed Killer controls most other broadleaf weeds, including sensitive weeds such as chickweed, clover, dandelion, knotweed and plantain.

OTHER USE PRECAUTIONS:

Do not allow drift onto vegetables, flowers, ornamental plants, shrubs, trees and other desirable plants since injury may result. Do not pour spray solutions near these plants.

Whether using a hose end sprayer or pressure sprayer, do not use sprays that are fine because they can drift and cause damage to nontarget plants. Coarse droplet sprays are less likely to drift. Use only lawn-type sprayers.

Do not spray on carpetgrass, dichondra, Lippia, nor on lawns or turf where desirable clovers are present. Do not spray roots of trees and ornamentals.

Do not exceed specified dosages for any area; be particularly careful within dripline of tree and other ornamental species.

This product may be injurious to St. Augustinegrass and centipedegrass if misapplications occur. Do not apply this product to Floratam® St. Augustinegrass in Florida.

The maximum application rate to turf is 2.0 pounds 2,4-D acid equivalent per acre per application per site. The maximum number of broadcast applications per treatment site is 2 per year.

WHEN TO USE:

1. Spray at any time during the growing season when weeds are actively growing.
2. On new lawns, wait until the grass has reached a height of 2 inches before making an overall spray treatment. However, a spot treatment may be made at any time with a small pressure sprayer. Wait 3 weeks after treatment before seeding grass seed or planting sprigs or stolons.
3. Poor weed control may result if spray is applied during drought or just before rain.
4. For best results, mow before treatment.
5. Do not water lawn within 24 hours after treatment.

HOSE END SPRAYERS

HOW TO APPLY:

Shake well before applying. Connect spray nozzle to hose. Remove bottle cap and screw bottle onto nozzle. Turn the "ON/OFF" switch on top of the nozzle right to left to "OFF". Turn on the water at the tap. When ready to treat, turn the "ON/OFF" switch to its center "ON" position. The water mixes automatically with the product at the proper ratio. Walk and spray at a steady pace. Each mark on the side of bottle covers 1,000 to 1,200 square feet.

Use a hose end sprayer designed for home lawn care. Refer to Charts 1 and 2 for the proper amounts of Trimec® Ready Spray Lawn Weed Killer. Determine the area of turf to be treated. Pour the recommended amount of product (Column 1) into the bottle, add water to the indicated level (Column 2), and spray uniformly on turf area shown in Column 3.

CHART 1: Amounts of Trimec® Ready Spray Lawn Weed Killer for bahiagrass, bermudagrass, Kentucky bluegrass, fescues, perennial ryegrass, and zoysiagrass lawns applied with hose end sprayers.

Column 1 Amount of Lawn Weed Killer to Use		Column 2	Column 3
(Tablespoons)	(Fluid Ounces)	Fill Bottle to, Gallons	Area to Treat, Square Feet
10 to 12	5 to 6	5	950
20 to 24	10 to 12	10	1,900
30 to 36	15 to 18	15	2,850
40 to 48	20 to 24	20	3,800

Note: For these grasses apply the entire contents of this container (32 fluid ounces) on 5,000 to 6,000 square feet of turf.

CHART 2: Amounts of Trimec® Ready Spray Lawn Weed Killer for centipedegrass and St. Augustinegrass lawns applied with hose end sprayers.

Column 1 Amount of Lawn Weed Killer to Use		Column 2	Column 3
(Tablespoons)	(Fluid Ounces)	Fill Bottle to, Gallons	Area to Treat, Square Feet
6	3	5	950
12	6	10	1,900
18	9	15	2,850
24	12	20	3,800

Note: For St. Augustinegrass and centipedegrass apply the entire contents of this container (32 fluid ounces) on 10,000 square feet of turf.

**PRESSURE SPRAYERS, BACKPACK SPRAYERS, OR
OTHER MANUALLY OPERATED SPRAYERS:**

Prepare the appropriate amounts of spray mixtures according to Chart 3. Measure the recommended amount of Trimec® Ready Spray Lawn Weed Killer into the spray tank and dilute with the proper amount of water. Use a spray pattern with small droplets and treat the area indicated in Chart 3.

Chart 3.

A. For bahiagrass, bermudagrass, Kentucky bluegrass, fescues, perennial ryegrass, and zoysiagrass lawns.			
Amount of Lawn Weed Killer to Use		Amount of Water in Sprayer, Gallons	Area to Treat, Square Feet
(Tablespoons)	(Fluid Ounces)		
2 to 2.4	1 to 1.2	¾	190
4 to 4.8	2 to 2.4	1½	380
8 to 9.6	4 to 4.8	3	760

B. For centipedegrass and St. Augustinegrass lawns.			
Amount of Lawn Weed Killer to Use		Amount of Water in Sprayer, Gallons	Area to Treat, Square Feet
(Tablespoons)	(Fluid Ounces)		
2 to 2.4	1 to 1.2	¾	380
4 to 4.8	2 to 2.4	1½	760
8 to 9.6	4 to 4.8	3	1,520

SPOT SPRAYING:

To spray individual weeds, do not use a hose end sprayer. Mix 1 fluid ounce (2 Tablespoons) of Trimec® Ready Spray Lawn Weed Killer in ½ gallon of water in a pressure sprayer. Spray to wet individual weeds.

GENERAL BRUSH CONTROL:

For control of brush along roads and other similar noncrop areas, mix 10 to 14 fluid ounces of Trimec® Ready Spray Lawn Weed Killer in 3 gallons of water and apply over 1,000 square feet. Use the higher rate of product for larger plants or more dense brush stands. Refer to Table 1 for additional instructions for spray preparations with water.

Table 1. Mixing instructions for preparing 1 to 9 gallons of spray solution with water for foliar application.		
Spray Solution Desired, Gallons	Amount of Trimec® Ready Spray Lawn Weed Killer Required	
	Normal Rate	High Rate (For larger plants or more dense brush)
0.25 (32 fl.oz.)	0.83 fl.oz.	1.16 fl.oz.
1	3⅓ fl.oz.	4⅔ fl.oz.
2	6⅔ fl.oz.	9⅓ fl.oz.
3	10 fl.oz.	14 fl.oz.
4	13⅓ fl.oz.	18⅔ fl.oz.
5	16⅔ fl.oz.	23⅓ fl.oz.
6	20 fl.oz.	28 fl.oz.
7	23⅓ fl.oz.	32⅔ fl.oz.
8	26⅔ fl.oz.	37⅓ fl.oz.
9	30 fl.oz.	42 fl.oz.

Equal measures: 1.0 fl.oz. = 2 tablespoons of product
1.0 quart = 32 fl.oz. = 2 pints = 4 cups

Best results are obtained when brush is young and actively growing. Do not cut brush until the herbicide has translocated throughout the plant causing root death. Keep out of lakes, streams and ponds.

CONTROLS:

BRUSH AND VINES:

Alder	Elderberry	Olive, Russian
Ash	Elm	Peppervine
Aspen	Giant ragweed	Persimmon
Bindweed	Gooseberry	Pine
Birch	Hackberry	Pine, shortleaf
Black cherry	Hazel	Poison ivy
Black locust	Hemp	Poison oak
Blackberry	Himalayan blackberry	Poplar
Brambles	Honeylocust	Raspberr
Buckbrush	Honeysuckle	Redvine
Buffalo berry	Honeysuckle, Japanese	Sagebrush
Bush morningglory	Kudzu	Saltbush
Buttonbush	Lilac, common	Serviceberry
Cedar	Marijuana	Skunkbrush
Chamise	Manzanita	Snowberry
Cherry, wild	Matrimony vine	Spruce
Coastal sage	Moonseed vine	Sumac
Cottonwood	Mountain ash	Sumac, dwarf black
Cucumber tree	Multiflora rose	Sycamore
Dogwood	Oak	Tamarisk

Thistle
Trumpet Vine
Trumpet creeper

Virginia creeper
Wild plum
Willow

Winterberry & other brush, trees & vines

HARDY WEEDS:

For hardy weeds such as clover, ground ivy, oxalis, sheep sorrel, dock, thistles, and morningglory, and Virginia buttonweed in lawns and other areas, repeat application in two to three weeks if necessary.

CONTROLS:

SENSITIVE WEEDS:

Bedstraw
Black medic
Buckhorn
Chickweed
Clover

Dandelion
Knotweed
Lambsquarters
Peppergrass
Plantain

Purslane
Ragweed
Shepherdspurse
Wild lettuce

HARD-TO-KILL WEEDS:

Dock
Ground ivy
Healall
Lespedeza
Mallow
Morningglory

Oxalis
Pigweed
Poison ivy
Poison oak
Sheep sorrel
Speedwell

Spurge
Virginia buttonweed
Wild carrot
Wild onion
Yarrow & many other broadleaf weeds

OTHER WEEDS CONTROLLED:

Aster
Austrian fieldcress
Bedstraw
Beggarticks
Betony, Florida
Bindweed, field
Bird vetch
Bitter wintercress
Bittercress, hairy
Bitterweed
Black-eyed Susan
Black medic
Black mustard
Blackseed plantain
Blessed thistle
Bloodflower milkweed
Blue lettuce
Blue vervain
Bracted plantain
Bristly oxtingue
Broadleaf dock
Broadleaf plantain
Broomweed
Buckhorn plantain
Bulbous buttercup
Bull thistle
Bullnettle
Burclover
Burdock
Burning nettle
Burweed
Buttercup
Buttonweed
Canada thistle
Carolina geranium
Carpetweed
Catchweed bedstraw
Catnip
Catsear
Chickweed, common
Chickweed, mouseear
Chicory
Cinquefoil
Clover, crimson
Clover, hop
Clover, red
Clover, strawberry
Cover, sweet
Clover, white
Cockle
Cocklebur
Common mullein
Creeping jenny
Cudweed
Curly dock
Daisy, English
Daisy, fleabane
Daisy, oxeye
Dandelion
Dichondra
Dogfennel
Dollarweed
False dandelion

Falseflax
False sunflower
Fiddleneck
Florida pusley
Frenchweed
Galinsoga
Goldenrod
Ground ivy
Gumweed
Hairy fleabane
Hawkweed
Healall
Heartleaf drymary
Heath aster
Hedge bindweed
Hedge mustard
Henbit
Hoary cress
Hoary plantain
Hoary vervain
Horsenettle
Jimsonweed
Knawel
Knotweed
Kochia
Lambsquarters
Lespedeza
Mallow
Matchweed
Mexicanweed
Milk vetch
Morningglory
Mouseear hawkweed
Mugwort
Musk thistle
Mustard
Narrowleaf plantain
Narrowleaf vetch
Nettle
Orange hawkweed
Oriental cocklebur
Oxalis
Parsley-piert
Parsnip
Pearlwort
Pennycress
Pennywort
Pepperweed
Pigweed
Pineywoods bedstraw
Plains coreopsis
Plantain
Poison ivy
Poison oak
Pokeweed
Poorjoe
Prairie sunflower
Prickly lettuce
Prickly sida
Prostrate knotweed
Prostrate pigweed
Prostrate spurge
Prostrate vervain

Puncturevine
Purslane, common
Ragweed
Red sorrel
Redroot pigweed
Redstem filaree
Rough cinquefoil
Rough fleabane
Russian pigweed
Russian thistle
Scarlet pimpernel
Scotch thistle
Sheep sorrel
Shepherdspurse
Slender plantain
Smallflower galinsoga
Smooth dock
Smooth pigweed
Sorrel
Sowthistle
Spanishneedles
Speedwell
Spiny amaranth
Spiny cocklebur
Spiny sowthistle
Spotted catsear
Spotted spurge
Spurweed
Stinging nettle
Strawberry, India mock
Tall nettle
Tall vervain
Tansy ragwort
Tansy mustard
Tanweed
Thistle
Trailing crownvetch
Tumble mustard
Tumble pigweed
Velvetleaf
Venice mallow
Virginia buttonweed
Virginia creeper
Virginia pepperweed
Wavyleaf bullthistle
Western clematis
Western salsify
White mustard
Wild aster
Wild buckwheat
Wild carrot
Wild four-o'clock
Wild garlic
Wild geranium
Wild lettuce
Wild marigold
Wild mustard
Wild onion
Wild parsnip
Wild radish
Wild rape
Wild strawberry
Wild sweet potato

Wild vetch
Woodsorrel
Woolly croton
Woolly morningglory
Woolly plantain

Wormseed
Yarrow
Yellowflower
pepperweed

Yellow rocket and other
broadleaf weeds

LIMITED WARRANTY AND DISCLAIMER

The manufacturer warrants only that the chemical composition of this product conforms to the ingredient statement given on the label, and that the product is reasonably suited for the labeled use when applied according to the Directions for Use.

THE MANUFACTURER NEITHER MAKES NOR INTENDS ANY OTHER EXPRESS OR IMPLIED WARRANTIES, INCLUDING ANY WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE, WHICH ARE EXPRESSLY DISCLAIMED. This limited warranty does not extend to the use of the product inconsistent with label instructions, warnings or cautions, or to use of the product under abnormal conditions such as drought, excessive rainfall, tornadoes, hurricanes, etc. These factors are beyond the control of the manufacturer or the seller. Any damages arising from a breach of the manufacturer's warranty shall be limited to direct damages, and shall not include indirect or consequential damages such as loss of profits or values, except as otherwise provided by law.

The terms of this Limited Warranty and Disclaimer cannot be varied by any written or verbal statements or agreements. No employee or agent of the seller is authorized to vary or exceed the terms of this Limited Warranty and Disclaimer in any manner.

827/6-2007 AP100906
EPA REG. NO. 2217-540

Manufactured By

An Employee-Owned Company
1217 West 12th Street
Kansas City, Missouri 64101
consumer.pbigordon.com

SPECIMEN
LABEL