

[CLICK HERE TO JUMP TO USE DIRECTIONS](#)

**GORDON'S®**  
**TRIMEC®**  
**READY SPRAY**  
**LAWN WEED KILLER**

**JUST CONNECT TO GARDEN HOSE  
 AND SPRAY! COVERS UP TO  
 5,000 SQ. FT. OF NORTHERN GRASSES.**

**ACTIVE INGREDIENTS:**

2,4-D, dimethylamine salt (CAS 2008-39-1) . . . . .	5.56%
Mecoprop-p, dimethylamine salt (CAS 66423-09-4) . . . . .	1.34%
Dicamba, dimethylamine salt (CAS 2300-66-5) . . . . .	0.62%
<b>INERT INGREDIENTS:</b> . . . . .	<b>92.48%</b>

TOTAL 100.00%

**THIS PRODUCT CONTAINS:**

- 4.62% 2,4-dichlorophenoxyacetic acid equivalent.
- 1.11% (+)-(R)-2-(2-methyl-4-chlorophenoxy)propionic acid equivalent.
- 0.51% 3,6-dichloro-o-anisic acid equivalent.
- Isomer Specific by AOAC Method.

**KEEP OUT OF REACH OF CHILDREN  
 CAUTION**



**READ THE ENTIRE LABEL FIRST.  
 OBSERVE ALL PRECAUTIONS AND  
 FOLLOW DIRECTIONS CAREFULLY.**

**PRECAUTIONARY STATEMENTS**

**Hazards to Humans and Domestic Animals**

**CAUTION:** Causes moderate eye irritation. Harmful if absorbed through skin or swallowed. Avoid contact with skin, eyes or clothing. Wear protective eyewear. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet. Remove and wash contaminated clothing before reuse.

First Aid	
<b>If in eyes:</b>	<ul style="list-style-type: none"> <li>• Hold eye open and rinse slowly and gently with water for 15-20 minutes.</li> <li>• Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.</li> <li>• Call a poison control center or doctor for treatment advice.</li> </ul>
<b>If swallowed:</b>	<ul style="list-style-type: none"> <li>• Call a poison control center or doctor immediately for treatment advice.</li> <li>• Have person sip a glass of water if able to swallow.</li> <li>• Do not induce vomiting unless told to do so by a poison control center or doctor.</li> <li>• Do not give anything by mouth to an unconscious person.</li> </ul>
<b>If on skin or clothing:</b>	<ul style="list-style-type: none"> <li>• Take off contaminated clothing.</li> <li>• Rinse skin immediately with plenty of water for 15-20 minutes.</li> <li>• Call a poison control center or doctor for treatment advice.</li> </ul> <p style="text-align: right;"><i>(cont. on next column)</i></p>

**First Aid (cont.)**

<b>If inhaled:</b>	<ul style="list-style-type: none"> <li>• Move person to fresh air.</li> <li>• If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth, if possible.</li> <li>• Call a poison control center or doctor for further treatment advice.</li> </ul>
--------------------	--

Have the product container or label with you when calling a poison control center or doctor or going for treatment. You may also call toll-free 1-877-800-5556 for emergency medical treatment information.

**Environmental Hazards**

This pesticide is toxic to fish and aquatic invertebrates and may adversely affect non-target plants. Do not apply directly to water. Do not contaminate water when disposing of equipment washwaters or rinsate.

This chemical has properties and characteristics associated with chemicals detected in groundwater. The use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination. Application around a cistern or well may result in contamination of drinking water or groundwater.

**DIRECTIONS FOR USE**

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

**1. GENERAL PRECAUTIONS AND RESTRICTIONS:**

- Do not apply this product in a way that will contact any person or pet, either directly or through drift. Keep people and pets out of the area during application.
- Do not allow people or pets to enter the treated area until sprays have dried.

**IMPORTANT:** Do not exceed specified dosages for any area.

**2. FOR USE ON LAWNS:**

This product is designed for use through this sprayer for controlling broadleaf weeds in bahiagrass, bermudagrass, Kentucky bluegrass, fescue, perennial ryegrass, zoysiagrass, centipedegrass, and St. Augustinegrass lawns, vacant lots, playgrounds and other similar areas.

For control of dandelion, chickweed, plantain, clover, and other listed weeds, apply when weeds are actively growing. Spray weeds such as wild garlic and wild onion when they appear in late fall and again in early spring.

Water lawn thoroughly 1 to 2 days before application. Do not water for 24 hours after application. Do not mow for one week before treating.

Spray entire lawn, wetting weeds and lawn grasses, do not saturate. Adjust water pressure at faucet to give coarse spray droplets. Do not apply when temperatures are expected to exceed 85°F within 24 hours following application.

Temporary injury may occur on bentgrass, bermudagrass, and red fescue lawn grasses. Do not apply to carpetgrass, dichondra, Lippia lawns, and lawns where desirable clovers are present.

Do not apply this product to improved varieties of 'Floritam' St. Augustinegrass in Florida.

Newly seeded lawns may be treated after they have been mowed at least three times.

**Other Use Precautions:**

- Do not apply as a fine mist because of potential for injury to desirable plants.
- Do not use this product on or near desirable plants, including contact of spray on exposed root systems or adventitious shoots within the drip line of desirable trees and shrubs, since injury may result.
- Do not apply directly to or near water, storm drains, gutters, sewers, or drainage ditches. Do not apply within 25 feet of rivers, fish ponds, lakes, streams, reservoirs, marshes, estuaries, bays, and oceans. Do not apply when windy. To prevent product run-off, do not over water the treated area(s) to the point of runoff or apply when raining or when rain is expected that day.

**3. HOW TO USE THE TRIMEC® READY-SPRAY**

Shake well before using. Connect attached spray nozzle to garden hose. Turn the water control "ON/OFF" valve on top of spray nozzle to the "OFF" position. Turn on water at faucet. Extend hose to the farthest

area of lawn to be treated and work back towards faucet so that you do not walk over treated areas. Using a coin or key, rotate the small product control valve located near front of nozzle forward to the "OPEN" position.

To begin spraying, point spray nozzle towards lawn and turn the water control "ON/OFF" valve to the "ON" position. The water automatically mixes with the product. Walk at a steady pace while spraying using an even sweeping motion, slightly overlapping treated areas. Refer to graduated scale on side of container to guide coverage.

To stop spraying, turn water control "ON/OFF" valve to the "OFF" position. Rotate small product control valve back to "CLOSED" position. Turn off water at faucet. To relieve pressure, turn water control "ON/OFF" valve to the "ON" position and point sprayer away from any desirable plants before removing unit from hose.

One quart of Gordon's® Trimec® Ready Spray Lawn Weed Killer will cover 5,000 square feet of lawn for bahiagrass, Kentucky bluegrass, bermudagrass, fescue, perennial ryegrass, and zoysiagrass.

For centipedegrass and St. Augustinegrass, one quart of Gordon's Trimec Ready Spray Lawn Weed Killer will treat 8,000 square feet of lawn.

**Limitations on broadcast applications to residential turfgrass:**

The maximum application rate is 6.4 fl. oz. of product per 1,000 sq. ft. per application. The maximum number of broadcast applications is limited to 2 per year with a minimum of 30 days between applications. The maximum seasonal rate is 12.8 fl. oz. of product per 1,000 sq. ft.

**4. WEEDS CONTROLLED:**

This product is effective on hard to control broadleaf weed species such as dandelion, white clover, sheep sorrel and spurge. In addition, Gordon's Trimec® Ready Spray Lawn Weed Killer controls susceptible weeds including chickweed, henbit, and plantain. A partial list is presented as follows:

**CONTROLS:**

**Sensitive Weeds:** Bedstraw, Black medic, Buckhorn, Chickweed, Clover, Dandelion, Knotweed, Lambsquarters, Peppergrass, Plantain, Purslane, Ragweed, Sheperdspurse, Wild lettuce.

**Hard-To-Kill Weeds:** Chickory, Dock, Ground ivy, Healall, Lespedeza, Mallow, Morningglory, Oxalis, Pigweed, Poison ivy, Poison oak, Sheep sorrel, Speedwell, Spurge, Virginia buttonweed, Wild carrot, Wild onion, yarrow and many other broadleaf weeds.

<b>WEEDS</b>	
Annual yellow sweetclover	Canada thistle
Aster	Carolina geranium
Austrian fieldcress	Carpetweed
Bedstraw	Catchweed bedstraw
Beggarticks	Catnip
Betony, Florida	Catsear
Bindweed, field	Chickweed, common
Bird vetch	Chickweed, mouseear
Bitter wintercress	Chicory
Bittercress, hairy	Cinquefoil
Bitterweed	Clover, crimson
Black-eyed Susan	Clover, hop
Black medic	Clover, red
Black mustard	Clover, strawberry
Blackseed plantain	Clover, sweet
Blessed thistle	Clover, white
Bloodflower milkweed	Cockle
Blue lettuce	Cocklebur
Blue vervain	Common mullein
Bracted plantain	Creeping jenny
Bristly oxtongue	Cudweed
Broadleaf dock	Curly dock
Broadleaf plantain	Daisy, English
Broomweed	Daisy, oxeye
Buckhorn	Dandelion
Buckhorn plantain	Dichondra
Bulbous buttercup	Dogfennel
Bull thistle	Dollarweed
Bullnettle	False dandelion
Burclover	Falseflax
Burdock	False sunflower
Burning nettle	Fiddleneck
Burweed	Florida pusley
Buttercup	
Buttonweed	

(cont. on next column)

**WEEDS (cont.)**

Frenchweed	Russian pigweed
Galinsoga	Russian thistle
Goldenrod	Scarlet pimpernel
Ground ivy	Scotch thistle
Gumweed	Sheep sorrel
Hairy fleabane	Sheperdspurse
Hawkweed	Slender plantain
Healall	Smallflower galinsoga
Heartleaf drymary	Smooth dock
Heath aster	Smooth pigweed
Hedge bindweed	Sorrel
Hedge mustard	Sowthistle
Henbit	Spanishneedles
Hoary cress	Speedwell
Hoary plantain	Spiny amaranth
Hoary vervain	Spiny cocklebur
Horsenettle	Spiny sowthistle
Jimsonweed	Spotted catsear
Knawel	Spotted spurge
Knotweed	Spurweed
Kochia	Stinging nettle
Lambsquarters	Strawberry, India mock
Lespedeza	Tall nettle
Mallow	Tall vervain
Matchweed	Tansy ragwort
Mexicanweed	Tansy mustard
Milk vetch	Tanweed
Morningglory	Thistle
Mouseear hawkweed	Trailing crownvetch
Mugwort	Tumble mustard
Musk thistle	Tumble pigweed
Mustard	Velvetleaf
Narrowleaf plantain	Venice mallow
Narrowleaf vetch	Virginia buttonweed
Nettle	Virginia creeper
Orange hawkweed	Virginia pepperweed
Oriental cocklebur	Wavyleaf bullthistle
Oxalis	Western clematis
Parsley-piert	Western salsify
Parsnip	White mustard
Pearwort	Wild aster
Pennycress	Wild buckwheat
Pepperweed	Wild carrot
Pigweed	Wild four-o'clock
Pineywoods bedstraw	Wild garlic
Plains coreopsis	Wild geranium
Plantain	Wild lettuce
Poison ivy	Wild marigold
Poison oak	Wild mustard
Pokeweed	Wild onion
Poorjoe	Wild parsnip
Prairie sunflower	Wild radish
Prickly lettuce	Wild rape
Prickly sida	Wild strawberry
Prostrate knotweed	Wild sweet potato
Prostrate pigweed	Wild vetch
Prostrate spurge	Woodsorrel
Prostrate vervain	Woolly croton
Puncturevine	Woolly morningglory
Purslane, common	Woolly plantain
Ragweed	Wormseed
Red sorrel	Yarrow
Redroot pigweed	Yellow rocket
Redstem filaree	Yellowflower pepperweed
Rough cinquefoil	and other broadleaf weeds
Rough fleabane	

## STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage and disposal.

**PESTICIDE STORAGE:** Keep from freezing. Store in original container and place in locked storage area when not using.

**PESTICIDE DISPOSAL AND CONTAINER HANDLING:** Nonrefillable container. Do not reuse or refill this container. **If empty:** Place in trash or offer for recycling if available. **If partly filled:** Call you local solid waste agency for disposal instructions. Never place unused product down any indoor or outdoor drain.

## LIMITED WARRANTY AND DISCLAIMER

FOR USE ONLY AS DIRECTED.

TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THE MANUFACTURER NEITHER MAKES NOR INTENDS ANY EXPRESS OR IMPLIED WARRANTIES, INCLUDING ANY WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE, WHICH ARE HEREBY EXPRESSLY DISCLAIMED. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, IN NO CASE SHALL THE MANUFACTURER BE LIABLE FOR INCIDENTAL, CONSEQUENTIAL, OR SPECIAL DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT. If these terms are not acceptable, return this product unopened immediately to the point of purchase, and the purchase price will be refunded in full. The terms of this LIMITED WARRANTY AND DISCLAIMER cannot be varied by any written or verbal statements or agreements at the point of sale or elsewhere.

827/2-2010 AP060309  
EPA REG. NO. 2217-540



*An Employee-Owned Company*

MANUFACTURED BY  
PBI/GORDON CORPORATION  
1217 WEST 12TH STREET  
KANSAS CITY, MISSOURI 64101  
[www.pbigordon.com](http://www.pbigordon.com)

SPECIMEN  
LABEL