

[CLICK HERE TO JUMP TO USE DIRECTIONS](#)

**GORDON'S®**

# TRIMEC®

## WEED & FEED 24-4-8

**CONTROLS DANDELION, CHICKWEED, KNOTWEED, PLANTAIN, HENBIT, SPURGE and many other broadleaf weeds including those listed on this page.**

**ACTIVE INGREDIENTS:**

2,4-dichlorophenoxyacetic acid . . . . .	0.62%
(+)-(R)-2-(2-methyl-4-chlorophenoxy) propionic acid . . . . .	0.14%
Dicamba 3,6-dichloro-o-anisic acid . . . . .	0.06%
<b>INERT INGREDIENTS:</b> . . . . .	<b>99.18%</b>
<b>TOTAL 100.00%</b>	

**THIS PRODUCT CONTAINS:**

12.4 lbs. 2,4-dichlorophenoxyacetic acid equivalent per ton.  
 2.8 lbs. (+)-(R)-2-(2-methyl-4-chlorophenoxy) propionic acid equivalent per ton.  
 1.2 lbs. 3,6-dichloro-o-anisic acid equivalent per ton.  
 Isomer Specific by AOAC Methods.  
 Contains the single isomer form of Mecoprop-p.  
 TRIMEC® is a registered trademark of PBI/GORDON CORPORATION.

## KEEP OUT OF REACH OF CHILDREN CAUTION

See below for additional Precautionary Statements and First Aid.

**GUARANTEED ANALYSIS 24-04-08**

Total Nitrogen (N) . . . . .	24.00%
1.57% Ammoniacal Nitrogen	
22.43% Urea	
Available Phosphate (P <sub>2</sub> O <sub>5</sub> ) . . . . .	4.00%
Soluble Potash (K <sub>2</sub> O) . . . . .	8.00%
Derived From: Diammonium Phosphate, Muriate of Potash and Urea Nitrogen.	
Equivalent To: 538 Lbs. CaCO <sub>3</sub>	



**READ THE ENTIRE LABEL FIRST. OBSERVE ALL PRECAUTIONS AND FOLLOW DIRECTIONS CAREFULLY.**

### PRECAUTIONARY STATEMENTS

**Hazards to Humans and Domestic Animals**

**CAUTION:** Harmful if swallowed. Do not inhale dust. Do not allow contact with skin, eyes or clothing. When using this product, wear long-sleeved shirt, long pants, socks and shoes.

After using this product, remove clothing and launder separately before reuse, and promptly and thoroughly wash hands and exposed skin with soap and water.

### FIRST AID

<b>If in eyes:</b>	<ul style="list-style-type: none"> <li>Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.</li> <li>Call a poison control center or doctor for treatment advice.</li> </ul>
<b>If on skin or clothing:</b>	<ul style="list-style-type: none"> <li>Take off contaminated clothing.</li> <li>Rinse skin immediately with plenty of water for 15-20 minutes.</li> <li>Call a poison control center or doctor for treatment advice.</li> </ul>
<b>If swallowed:</b>	<ul style="list-style-type: none"> <li>Call a poison control center or doctor immediately for treatment advice.</li> <li>Have person sip a glass of water if able to swallow.</li> <li>Do not induce vomiting unless told to do so by a poison control center or doctor.</li> <li>Do not give anything by mouth to an unconscious person.</li> </ul>

*(Continued on next column)*

### FIRST AID (continued)

<b>If inhaled:</b>	<ul style="list-style-type: none"> <li>Move person to fresh air.</li> <li>If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth if possible.</li> <li>Call a poison control center or doctor for treatment advice.</li> </ul>
Have the product container or label with you when calling a poison control center or doctor or going for treatment. For additional information in case of emergency, call toll free 1-877-800-5556.	

**Environmental Hazards:** Drift or runoff may adversely affect nontarget plants. Do not apply directly to water. Do not contaminate water when disposing of equipment washwater. Do not apply when weather conditions favor drift from target area.

### DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Do not allow people (other than applicator) or pets on treatment area during application. Do not enter treatment area until spray has dried or dust has settled.

### STORAGE AND DISPOSAL

**STORAGE:** Store in original container and keep any bags containing unused product in a locked storage area inaccessible to children and domestic animals. Do not store near food or feedstuffs.

**CONTAINER DISPOSAL:** If empty: Do not reuse this container. Place in trash or offer for recycling if available. If partially filled: Call your local solid waste agency or toll free number (1-800-CLEANUP) for disposal instructions. Never place unused product down any indoor or outdoor drain.

### USE PRECAUTIONS

Do not use this product where desirable clovers are present or allow the material to drift onto flowers, vegetables or ornamental shrubbery. Do not apply in areas underlaid by roots of desirable trees or shrubs. Apply when the air is calm. Treated areas may be seeded 3 to 4 weeks after application. Application of this product to newly seeded areas should be delayed until the third mowing. During application, close spreader openings when turning.


Do not apply this product to dichondra or carpetgrass. Do not apply this product to 'Floritam' St. Augustinegrass in Florida.

Keep this product off sidewalks, patios, and driveways. This fertilizer may stain concrete and painted surfaces. If it comes in contact with concrete surfaces, sweep off immediately before flushing with water. If concrete is wet when contact occurs, staining may be impossible to prevent. Be careful when using rotary spreader which scatter fertilizer beyond lawn edge and onto concrete.


### CALIBRATION INSTRUCTIONS

To provide proper distribution, calibrate spreader before application.


- Always apply with a properly calibrated rotary or drop fertilizer spreader to distribute the granules evenly. Never apply by hand.
- Place your spreader on the driveway or other paved surface when filling. This guards against spilling fertilizer on grass. For even coverage, begin walking before you turn spreader to the open position to avoid dropping fertilizer in one spot when the spreader is opened. Be sure the spreader is closed when filling, when turning to backtrack, or when stopping.




**1. MOW LAWN**  
to normal height 1 to 2 days before application.





**2. WATER LAWN**  
at least 1 day before application to sustain moisture until the next watering (see step 5).



**3. SPREADER SETTINGS\***  
 Scott . . . . . .6  
 Lawn Beauty . . . . .5  
 Jackson . . . . .7½  
 Sears . . . . . .4



**4. APPLICATION TO MOIST LAWN**  
Apply when weeds are young and actively growing, preferably in the morning when the dew is on the grass. If grass is not moist at time of application, sprinkle lightly with water to hold the material and prevent dust drift onto nontarget plants.



**5. WATER-IN THOROUGHLY**  
Do not wash from weed leaves for 1 to 2 days after application. At this time, a thorough watering should be made.

\*The above spreader settings are approximate. Calibrate your spreader on 275 square feet and adjust to apply at the rate of 1 lb. of product per 275 square feet.

## DISTRIBUTION AND SPREADER PATTERNS

**Coverage** – To determine how much you need to apply, multiply the length of your lawn by its width. This equals your total square footage. Deduct all non-lawn areas (house, gardens, etc.). One bag covers up to 5,000 square feet.

## SINGLE PASS DISTRIBUTION

First spread the borders of your lawn. Then spread the center by spreading in straight line passes. Spread at a uniform walking speed to insure even coverage. To avoid gaps and streaks, first spread the border and fill in center with uniform, parallel passes.

## INSTRUCTIONS

GORDON'S TRIMEC WEED & FEED is a Trimec® herbicide, a combination of three proven weed killers — 2,4-D, mecoprop-p and dicamba. Together, they control a wide range of lawn weeds including dandelion, chickweed, knotweed, plantain, henbit, spurge and many others commonly found in home lawns. In addition, a 24-4-8 fertilizer provides a supply of plant food. Retreatment may be necessary if (a) lawn is heavily overrun by weeds, or, (b) if adverse low moisture conditions prevail and weeds are in a state of poor growth. A second application may be made in approximately 30 days.

GORDON'S TRIMEC WEED & FEED controls broadleaf weeds commonly found in cool season and warm season turfgrass established in residential lawns. Established turf of Kentucky bluegrass, fescues, and perennial ryegrass may be treated. Established turf of bermudagrass, bahiagrass, buffalograss, zoysiagrass, St. Augustinegrass and centipedegrass may be treated.

The maximum application rate to turf is 1.0 pound of 2,4-D acid equivalent per acre per application per site. The maximum number of broadcast applications per treatment site is 2 per year.

This product has been registered with the Washington State Department of Agriculture. When applied as directed, this fertilizer meets the Washington standards for arsenic, cadmium, cobalt, mercury, molybdenum, lead, nickel, selenium, and zinc. You have the right to receive specific information about Washington standards from the distributor of this product. Information regarding the contents and levels of metals in this product is available on the internet at <http://www.aapco.org/metals.htm>.

## BROADLEAF WEEDS

Annual yellow sweetclover	Clover, red	Milk vetch
Aster	Clover, strawberry	Morningglory
Austrian fieldcress	Clover, sweet	Mouseear hawkweed
Bedstraw	Clover, white	Mugwort
Beggarticks	Cockle	Musk thistle
Betony, Florida	Cocklebur	Mustard
Bindweed, field	Common mullein	Narrowleaf plantain
Bird vetch	Creeping jenny	Narrowleaf vetch
Bitter wintercress	Cudweed	Nettle
Bittercress, hairy	Curly dock	Orange hawkweed
Bitterweed	Daisy, English	Oriental cocklebur
Black-eyed Susan	Daisy fleabane	Oxalis
Black medic	Daisy, oxeeye	Parsley-piert
Black mustard	Dandelion	Parsnip
Blackseed plantain	Dichondra	Pearlwort
Blessed thistle	Dogfennel	Pennycress
Bloodflower milkweed	Dollarweed	Pepperweed
Blue lettuce	False dandelion	Pigweed
Blue vervain	Falseflax	Pineywoods bedstraw
Bracted plantain	False sunflower	Plains coreopsis
Bristly oxtongue	Fiddleneck	Plantain
Broadleaf dock	Florida pusley	Poison ivy
Broadleaf plantain	Frenchweed	Poison oak
Broomweed	Galinsoga	Pokeweed
Buckhorn	Goldenrod	Poorjoe
Buckhorn plantain	Ground ivy	Prairie sunflower
Bulbous buttercup	Gumweed	Prickly lettuce
Bull thistle	Hairy fleabane	Prickly sida
Bullnettle	Hawkweed	Prostrate knotweed
Burclover	Healall	Prostrate pigweed
Burdock	Heartleaf drymary	Prostrate spurge
Burning nettle	Heathaster	Prostrate vervain
Burweed	Hedge bindweed	Puncturevine
Buttercup	Hedge mustard	Purslane, common
Buttonweed	Henbit	Ragweed
Canada thistle	Hoary cress	Red sorrel
Carolina geranium	Hoary plantain	Redroot pigweed
Carpetweed	Hoary vervain	Redstem filaree
Catchweed bedstraw	Horsenettle	Rough cinquefoil
Catnip	Jimsonweed	Rough fleabane
Catsear	Knawel	Russian pigweed
Chickweed, common	Knotweed	Russian thistle
Chickweed, mouseear	Kochia	Scarlet pimpernel
Chicory	Lambsquarters	Scotch thistle
Cinquefoil	Lespedeza	Sheep sorrel
Clover, crimson	Mallow	Shepherdspurge
Clover, hop	Matchweed	Slender plantain
	Mexicanweed	Smallflower galinsoga

Smooth dock	Trailing crownvetch	Wild mustard
Smooth pigweed	Tumble mustard	Wild onion
Sorrel	Tumble pigweed	Wild parsnip
Sowthistle	Velvetleaf	Wild radish
Spanishneedles	Venice mallow	Wild rape
Speedwell	Virginia buttonweed	Wild strawberry
Spiny amaranth	Virginia creeper	Wild sweet potato
Spiny cocklebur	Virginia pepperweed	Wild vetch
Spiny sowthistle	Wavyleaf bullthistle	Woodsorrel
Spotted catsear	Western clematis	Woolly croton
Spotted spurge	Western salsify	Woolly morningglory
Spurweed	White mustard	Woolly plantain
Stinging nettle	Wild aster	Wormseed
Strawberry, India mock	Wild buckwheat	Yarrow
Tall nettle	Wild carrot	Yellow rocket
Tall vervain	Wild four-o'clock	Yellowflower
Tansy ragwort	Wild garlic	pepperweed
Tansy mustard	Wild geranium	and other
Tanweed	Wild lettuce	broadleaf weeds
Thistle	Wild marigold	

## LIMITED WARRANTY AND DISCLAIMER

The manufacturer warrants only that the chemical composition of this product conforms to the ingredient statement given on the label, and that the product is reasonably suited for the labeled use when applied according to the Directions for Use.

**THE MANUFACTURER NEITHER MAKES NOR INTENDS ANY OTHER EXPRESS OR IMPLIED WARRANTIES, INCLUDING ANY WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE, WHICH ARE EXPRESSLY DISCLAIMED.** This limited warranty does not extend to the use of the product inconsistent with label instructions, warnings or cautions, or to use of the product under abnormal conditions such as drought, excessive rainfall, tornadoes, hurricanes, etc. These factors are beyond the control of the manufacturer or the seller. Any damages arising from a breach of the manufacturer's warranty shall be limited to direct damages, and shall not include indirect or consequential damages such as loss of profits or values, except as otherwise provided by law.

The terms of this Limited Warranty and Disclaimer cannot be varied by any written or verbal statements or agreements. No employee or agent of the seller is authorized to vary or exceed the terms of this Limited Warranty and Disclaimer in any manner.

570/7-2004 AP031804  
EPA REG. NO. 2217-658-33955

Manufactured and Distributed By  
 **pbi/gordon**  
**CORPORATION**  
 An Employee-Owned Company  
 1217 West 12th Street  
 Kansas City, Missouri 64101  
[www.pbigordon.com](http://www.pbigordon.com)